

Southeast Regional Muscadine Grape

INTEGRATED
MANAGEMENT
GUIDE

2022



UNIVERSITY OF GEORGIA
EXTENSION

2022 Southeast Regional Muscadine Grape Integrated Management Guide

Commodity Editor

Bill Cline, North Carolina State University

Senior Editors

Phil Brannen, University of Georgia

Bill Cline, North Carolina State University

Section Editors

Pathology

Bill Cline, North Carolina State University

Phil Brannen, University of Georgia

Rebecca Melanson, Mississippi State University

Mary Helen Ferguson, Louisiana State University

Entomology

Brett Blaauw, University of Georgia

Frank Hale, University of Tennessee

Meredith Favre, North Carolina State University

Aaron Cato, University of Arkansas

Weed Science

Wayne Mitchem, North Carolina State University

Vertebrate Management

David Lockwood, University of Tennessee

Michael T. Mengak, University of Georgia

Pesticide Stewardship and Safety

Ash Sial, University of Georgia

Recommendations are based on information from the manufacturer's label and performance data from research and extension field tests.

Because environmental conditions and grower application methods vary widely, suggested use does not imply that performance of the pesticide will always conform to the safety and pest control standards indicated by experimental data.

This publication is intended for use only as a guide. Specific rates and applications methods are on the pesticide label, and these are subject to change at any time. Registrations also vary between states and are subject to change at any time, please check with your state department of agriculture or regulatory agency concerning current registration status within your state. Always refer to and read the pesticide label before making any application! The pesticide label supersedes any information contained in this guide, and it is the legal document referenced for application standards.

TABLE OF CONTENTS

| | |
|---|-----------|
| Pesticide Emergencies (contacts) | 3 |
| Pesticide Liability and Stewardship | 5 |
| Integrated Management Guide (Insect and Disease Control) | 8 |
| Cultural Practices | 9 |
| Fungicide Efficacy Chart | 33 |
| Weed Management Guide | 35 |
| Herbicide Resistance Management Information | 35 |
| Suggested Herbicide Programs for Grape Vineyards | 42 |
| Wildlife Damage and Prevention | 46 |

Southeastern Region University Small Fruits IPM Contacts

| | | | | |
|--|---|---|--|---|
| <p>Auburn University Elina Coneva (<i>Horticulture</i>) Ed Sikora (<i>Pathology</i>)</p> <p>University of Arkansas Aaron Cato (<i>IPM</i>) Amanda McWhirt (<i>Horticulture</i>)</p> <p>Clemson University Wayne Mitchem (<i>Weed Science</i>) Guido Schnabel (<i>Pathology</i>)</p> | <p>University of Florida Phil Harmon (<i>Pathology</i>) Oscar Liburd (<i>Entomology</i>) Natalia Perez (<i>Pathology</i>) Jeff Williamson (<i>Horticulture</i>)</p> <p>University of Georgia Mark Czarnota (<i>Weed Science</i>) Brett Blaauw (<i>Entomology</i>) Phil Brannen (<i>Pathology</i>) Elizabeth Little (<i>Organic Pathology</i>) Michael T. Mengak (<i>Wildlife</i>) Wayne Mitchem (<i>Weed Science</i>) Jonathan Oliver (<i>Pathology</i>) Ash Sial (<i>Entomology</i>)</p> | <p>University of Kentucky Daniel Becker (<i>Horticulture</i>) Ric Bessin (<i>Entomology</i>) Nicole Gauthier (<i>Pathology</i>) John Strang (<i>Horticulture</i>) Patsy Wilson (<i>Grapes, Horticulture</i>) Shawn Wright (<i>Weeds, Horticulture</i>)</p> <p>Louisiana State University Mary Helen Ferguson (<i>Pathology</i>) Kathryn Fontenot (<i>Horticulture</i>) Raghuwinder Singh (<i>Pathology</i>) Ron Strahan (<i>Weed Science</i>)</p> | <p>Mississippi State University Blake Layton (<i>Entomology</i>) Rebecca A. Melanson (<i>Pathology</i>) Eric Stafne (<i>Horticulture</i>)</p> <p>North Carolina State University Ryan Adams (<i>IPM</i>) Bill Cline (<i>Pathology</i>) Gina Fernandez (<i>Horticulture</i>) Katie Jennings (<i>Weed Science</i>) Mark Hoffmann (<i>Horticulture</i>) Wayne Mitchem (<i>Weed Science</i>) Sara Villani (<i>Pathology</i>)</p> | <p>University of Tennessee Zach Hansen (<i>Pathology</i>) David Lockwood (<i>Horticulture</i>)</p> <p>Virginia Tech Jeff Derr (<i>Weed Science</i>) Mizuho Nita (<i>Pathology</i>) Doug Pfeiffer (<i>Entomology</i>) Jayesh Samtani (<i>Horticulture</i>)</p> |
|--|---|---|--|---|

PESTICIDE EMERGENCIES

1-800-222-1222

This number automatically connects you with a local Poison Control Center from anywhere in the United States.

Symptoms of Pesticide Exposure

- **Tightening of the chest, mental confusion, blurred vision, rapid pulse, intense thirst, vomiting, convulsions, and unconsciousness are always serious symptoms! Dial 911!**
- Pesticides with 'DANGER' or 'DANGER/POISON' on the product label can cause severe injuries or death very quickly, even with small exposures. **Take immediate action!**
- *Other symptoms of pesticide poisoning:* headache, fatigue, weakness, restlessness, nervousness, profuse sweating, tearing and drooling, nausea, diarrhea, or irritation of the skin/ eyes/nose/throat. Consult the product Material Safety Data Sheet (MSDS) for symptoms associated with a particular pesticide.

Pesticide on Skin

- **WASH, WASH, WASH!** Immediately wash pesticide from skin as thoroughly as possible with any available water that does not contain pesticides.
- Quickly remove protective clothing and any contaminated clothing.
- *Rewash* contaminated skin with soap and water as soon as possible.
- **If the victim experiences any symptom(s) of poisoning, get medical assistance immediately.** *Take the pesticide label with you,* but do not contaminate vehicles or expose others if you must take the container with you.

Pesticide in Eyes

- Rinse eye(s) gently with *clean* water for *at least* 15 minutes. Be careful of water temperature.
- **If eye remains irritated or vision is blurry after rinsing, get medical attention right away!** *Take the pesticide label with you.*

Pesticide in Mouth or Swallowed

- Provide / drink large amounts of water or milk to drink - *Do not give liquids to a person who is unconscious or convulsing!*
- Consult the label BEFORE vomiting is induced – the label may advise against inducing vomiting. Do not induce vomiting with emulsifiable concentrate (E, EC) formulations.
- *Do not induce vomiting if a person is unconscious or is convulsing!*
- **Seek medical attention.** *Take the pesticide label with you.*
- If the pesticide was not swallowed, *rinse mouth thoroughly with clean water.* If mouth is burned or irritated, consult a physician.

Pesticide Inhaled

- Move victim to fresh air immediately!
- Warn others in the area of the danger.
- Loosen tight clothing.
- Administer artificial respiration if necessary, but try to determine if the person also may have swallowed any pesticide - avoid any pesticide or vomit that may be around the victim's mouth.
- **Seek medical attention.** *Take the pesticide label with you.*

Heat Stress

- Move the victim to a cooler area, remove protective clothing, and pour cool water over the person.
- Give cool liquids to drink – *Do not give liquids to a person who is unconscious or convulsing!*
- **Pesticide poisoning may mimic heat illness! Get medical attention if the person is unconscious or if the person is not fully recovered within 15 minutes of cooling down and drinking liquids.**

Signal Words

- The pesticide signal word will appear on the pesticide label. It provides information about the acute risks of the pesticide to people.
 - **DANGER/POISON:** *Highly toxic* - less than a teaspoon can kill an adult.
 - **DANGER:** *Highly toxic* - pesticide can cause severe eye and/or skin injury.
 - **WARNING:** *Moderately toxic* - two tablespoons or less can kill an adult.
 - **CAUTION:** *Slightly toxic* - an ounce or more is required to kill an adult.

Understand that the signal word does *not* provide information about long term pesticide exposure risks (e.g., cancer) or allergic effects. Minimize your exposure to *all* pesticides. The signal word does *not* indicate environmental toxicity or other environmental effects.

PESTICIDE SPILLS OR OTHER EMERGENCIES

Spills on public roads (Usually call the state police/state highway patrol. In many cases, you can call CHEMTREC for assistance 24 hours a day at 1-800-424-9300 or 911.)

| STATE | AGENCY | PHONE NUMBER |
|-----------------------|--|--|
| Alabama | Alabama Highway Patrol Post Alabama Department of Environmental Management Alabama Emergency Management Agency | *HP (*47) for mobile phones (334) 271-7700 (205) 280-2200 |
| Arkansas | Arkansas Department of Emergency Management | 800-322-4012 |
| Georgia | Georgia State Patrol | Cell: call *GSP or 911 |
| Louisiana | LDAF Emergency Hotline | 855-452-5323 |
| Mississippi | Mississippi Emergency Management Agency | 1-800-222-6362 |
| North Carolina | Regional Response Team (RRT) For spills not on public roadways, contact the Pesticide Section of NCDA&CS | 911 or your RRT (919) 733-3556 or (800) 662-7956 during non-business hours |
| South Carolina | South Carolina Highway Patrol ----- South Carolina DHEC Emergency Response Section | Cell: call *HP ----- 888-481-0125 (Toll Free) |
| Tennessee | Tennessee Emergency Management Agency (TEMA) State Emergency Operations Center | 800-262-3300 |
| Virginia | Virginia Emergency Operations Center | 804-674-2400 |

Environmental emergencies (contamination of waterways, fish kills, bird kills, etc.)

| STATE | AGENCY | PHONE NUMBER |
|-----------------------|--|--|
| Alabama | Alabama Department of Environmental Management Alabama Department of Emergency Management Alabama Department of Conservation and Natural Resources | 334-271-7700 205-280-2200 334-242-3469 |
| Arkansas | Arkansas Department of Emergency Management | 800-322-4012 (Toll Free) |
| Georgia | Georgia Department of Natural Resources Response Team | 800-241-4113 (Toll Free) |
| Louisiana | LDAF Emergency Hotline | 855-452-5323 (Toll Free) |
| Mississippi | Mississippi Emergency Management Agency | 1-800-222-6362 |
| North Carolina | North Carolina Div. of Water Quality | 800-858-0368 (Toll Free) |
| South Carolina | South Carolina DHEC | 888-481-0125 (Toll Free) |
| Tennessee | Tennessee Wildlife Resources Agency | Region 1 West Tennessee: 800-372-3928 Region 2 Middle Tennessee: 800-624-7406 Region 3 Cumberland Plateau: 800-262-6704 Region 4 East Tennessee: 800-332-0900 |
| Virginia | Virginia Emergency Operations Center | 1-804-674-2400 |

PESTICIDE LIABILITY and STEWARDSHIP

Pesticide applicators, supervisors, and business owners may all face severe criminal and/or civil penalties if pesticides are misused – knowingly or accidentally.

UNDERSTANDING PESTICIDE LABELS

Federal and state laws require pesticide applicators to follow the directions on the pesticide label exactly. Do not exceed maximum label rates, apply a pesticide more frequently than stated on the label, or apply a pesticide to a site that is not indicated on the label. Labels change; review yours regularly.

Restricted Use Pesticides (RUP). These pesticides are clearly labeled “Restricted Use Pesticide” in a box at the top of the front label. Applicators purchasing, applying, or supervising the application of a RUP, must be certified or licensed through their state pesticide regulatory agency. Some states have mandatory licensing for certain pesticide use categories whether or not RUPs are applied.

Personal Protective Equipment (PPE). Anyone handling or applying pesticides must wear the PPE stated on the pesticide label. The Worker Protection Standard requires applicators to wear the label required PPE and requires agricultural employers to supply the label PPE and ensure that the PPE is worn correctly by applicator employees. Do not wear PPE items longer than it has been designed to protect you. Clean, maintain and properly store PPE. Do not store PPE with pesticides.

Reentry Interval (REI). The period immediately following the application of a pesticide during which unprotected workers should not enter a field.

Pre-Harvest Interval (PHI). The time between the last pesticide application and harvest of the treated crops.

EPA Worker Protection Standard (WPS) WPS changes continue to be implemented. Growers should consult the EPA website (<https://www.epa.gov/pesticide-worker-safety/agricultural-worker-protection-standard-wps>) or their local extension service for the most up to date information. Growers who employ one or more *non-family* members must comply with the WPS. This standard requires agricultural employers to protect applicator employees and agricultural worker employees

from pesticide exposure in the workplace by 1) providing specified pesticide safety training, 2) providing specific information about pesticide applications made on the agricultural operation, 3) providing and ensuring that applicators wear clean and properly maintained label required PPE, 4) providing decontamination facilities for potential pesticide and pesticide residue exposures, and 5) providing timely access to medical assistance in the event of a suspected pesticide exposure. These protections apply to both restricted use *and* general use pesticides used in agricultural plant production.

Pesticide Recordkeeping. You must keep records of all RUP applications for at least two years under the Federal (USDA) Pesticide Recordkeeping Requirement if your state does not have its own pesticide recordkeeping requirements. Some states require records be kept for longer than the federal requirement. Maintaining records of all pesticide applications, not just RUP applications, indefinitely, cannot only help troubleshoot application problems, but also allows you to reference successful applications and can help protect against future liability. Consult your [local Extension Service](#) for details.

Emergency Preparedness: Be prepared for emergencies. Store pesticides and clean empty containers securely. Develop and provide written plans and training to prepare your employees, and family members, for pesticide fires, spills, and other emergencies. Assign responsibilities to be carried out in the event of pesticide emergencies. Keep copies of the pesticide labels and MSDS away from the area where pesticides are stored. Provide copies of product MSDSs to your community first responders. Consult your [local Extension Service](#) and insurance company for assistance.

Pesticide Disposal: Properly dispose of clean empty pesticide containers and unwanted pesticides as soon as possible. Containers can often be recycled in a pesticide container recycling program. Unwanted pesticides may pose a risk of human exposure and environmental harm if kept for long periods of time. Consult [local Extension Service](#) for assistance.

PESTICIDE APPLICATION

Information on pesticide use is available from the Pesticide Environmental Stewardship website (<http://pesticidestewardship.org>) including information on [sprayer calibration \(airblast sprayer calibration can be found here\)](#), [personal protective equipment](#), [recordkeeping](#), and [resistance management](#).

ORGANICALLY ACCEPTABLE MATERIALS

Products that are listed by the Organic Materials Review Institute (OMRI, www.omri.org) are commonly accepted for use in certified organic production systems. However, always consult your organic certifier prior to using a product as certifiers may differ in what products they accept. **OMRI**-listed materials are indicated in the Comments section.

Mode of Action (MOA): Pesticides affect their target pest in a variety of ways, and the way a pesticide kills the target organism is called the *mode of action* (MOA). Although pesticides have different names and may have different active ingredients, they may have the same MOA. Over time, pests can become resistant to a pesticide, and typically this resistance applies to all pesticides with the same MOA. When rotating pesticides, it is important to select pesticides with different MOAs.

The **Fungicide Resistance Action Committee (FRAC)**, **Insecticide Resistance Action Committee (IRAC)** and IRAC Nematode Working Group, and the **Weed Science Society of America (WSSA)** have organized crop protection materials into groups with shared MOAs and given them specific codes, which appear on pesticide labels and are listed in this guide to aid in the development of resistance management programs. Some MOAs may be unknown and given a code with a U. *When selecting pesticides, avoid successive applications of materials in the same MOA group to minimize potential resistance development.* MOA categories are listed in this guide to aid in the development of resistance management programs. More information about this topic can be found at www.frac.info, www.irac-online.org, and www.hracglobal.com.

RESISTANCE MANAGEMENT

Insects, weeds, and disease-causing organisms are all capable of developing resistance to pesticides. To minimize the likelihood of resistance development against your material of choice:

1. Only use pesticides when necessary: When the damage caused by the pest you are controlling is greater than the cost of the pesticide and no other, effective options are available.
2. Use the appropriate material for the pest.
3. Use the recommended rate of the material. Do not use a lower rate than listed on the label.
4. If more than one treatment is needed when the same pest is present, rotate pesticide mode of action (MOA) between treatments.

State Registrations: This publication is a regional guide, and as such every product listed may not be available or registered for use in every state. We made every effort to indicate these materials in the Comments section, but because state level registration status may change, verify that a product is registered for use in your state before purchasing and applying it. This may be done by visiting one of several online databases (examples provided below) that provide information on the state registration status of various products, by visiting product manufacturer websites, or by contacting your local county Extension agent or an appropriate state Extension specialist.

| Database | Web Address |
|--|---|
| Agrian Label Database | https://home.agrian.com/ |
| Crop Data Management Systems | http://www.cdms.net/Label-Database |
| EPA Pesticide Product and Label System | https://iaspub.epa.gov/apex/pesticides/f?p=PPLS:1 |
| Greenbook Data Solutions | https://www.greenbook.net/ |
| Kelly Registration Systems ¹ | http://www.kellysolutions.com |
| National Pesticide Information Retrieval System ² | http://npirspublic.ceris.purdue.edu/state/ |

¹Available for AL, FL, GA, MS, NC, SC, and VA in the southeastern U.S.

²Available for AL, AR, FL, KY, LA, TX, VA, and WV in the southeastern U.S.

POLLINATOR PROTECTION

Before making insecticide applications, monitor insect populations to determine if treatment is needed. If pesticide (insecticide or fungicide) application is necessary:

1. Use selective pesticides to reduce risk to pollinators and other non-target beneficial insects.
2. Read and follow all pesticide label directions and precautions (the Label is the Law!). EPA now requires the addition of a “Protection of Pollinators” advisory box on certain pesticide labels. Look for the bee hazard icon in the Directions for Use and within crop specific sections for instructions to protect bees and other insect pollinators.



(Continued on next page)

3. Minimize in-field exposure of bees to pesticides by avoiding applications when bees are actively foraging in the crops. Bee flower visitation rate is highest in early morning. Apply pesticides in the late afternoon or early evening to allow for maximum residue degradation before bees return the next morning. Bee foraging activity is also dependent upon time of year (temperature) and stage of crop growth. The greatest risk of bee exposure is during bloom.
4. Minimize off-target movement of pesticide applications by following label directions to minimize drift. Do not make pesticide applications when the wind is blowing towards beehives or off-site pollinator habitats.

Muscadine Grape Integrated Management Guide

Introduction

The muscadine grape (*Vitis rotundifolia*, syn. *Muscadinia rotundifolia*) is a thick-skinned native grape species found in the southeastern United States. The fruit ripens in late summer/early fall and is generally harvested as single berries rather than in bunches. In the wild, muscadine vines are usually either male or female, while cultivated varieties (cultivars) are usually either female or perfect-flowered (having both male and female flower parts). Most wild vines have dark fruit, while cultivated (improved) cultivars are also available in many shades of bronze, purple or red. Cultivars have been developed for both fresh fruit and for winemaking. As the species name implies, the leaves are round and unlobed, though serrated at the edge. NOTE: For purposes of pesticide labeling, muscadines are included with all other grapes.

Diseases – Muscadines can be grown in warm, humid areas where bunch grapes cannot, and have been grown organically for local markets. Muscadines are resistant to the Pierce’s Disease bacterium (*Xylella fastidiosa*) that limits bunch grape production and are not affected by some major fungal diseases like grape downy mildew or botrytis bunch rot. However, muscadines can still suffer extensive losses if diseases like powdery mildew, bitter rot, black rot and ripe rot are left unchecked. Fruit rotting diseases reduce yields and make berries unmarketable for either wine or fresh sales. **Spray volume** – Muscadines form a dense canopy that is difficult to penetrate with spray solution. Directed nozzles and spray volumes of 100 gallons per acre or more may be needed to ensure adequate coverage. Test your sprayer by using water-sensitive spray indicator cards suspended in the fruiting zone of the cordon to determine whether spray coverage is adequate. **Nematodes** – The impact of nematodes on muscadines remains undetermined. A survey in NC and GA determined the prevalent species associated with muscadines but found no correlation with plant health. Muscadines are likely less susceptible than other grape species, however testing for nematodes is advisable when replanting old vineyard sites.

Insects – See text in each section below for specific comments. Grape root borers are an important pest of muscadine vines. Aphids occur on new shoots but often do not require control. Grape berry moth occurs sporadically. Japanese beetles may feed on foliage and flowers in early summer. Insects feeding on ripe fruit include green June beetles, as well as wasps, yellowjackets and other stinging insects that can pose a risk to pickers. The risk of infestation by spotted wing drosophila (SWD) is considered low due to the thick skins of muscadine grapes; however, SWD and other vinegar flies can be found in damaged fruit such as rain-splits.

Weeds—See comments beginning on page 24.

Cultural Practices

For comprehensive information on establishing vines and other muscadine production practices, see the *Muscadine Grape Production Guide for the Southeast*: <https://content.ces.ncsu.edu/muscadine-grape-production-guide>

Cultivar selection – Most muscadines grown for wine are high-yielding, medium to small size grapes that detach easily and thus are suitable for machine harvest. The cultivar Carlos is the most commonly used bronze muscadine for white wine, and Noble is the most common dark fruited cultivar for red wine. Fresh market and pick-your-own cultivars have grapes that are larger in size, with firmer flesh, and a more edible skin. Examples of fresh-market cultivars include dark-fruited Lane, Supreme, Paulk and Nesbitt, and the bronze-fruited cultivars Hall, Triumph, Early Fry, Summit, Fry and Late Fry.

Site selection and establishment – The most productive muscadine vineyards are on sites that have well-drained soils. Soils that stay wet during the winter months or that flood regularly are not satisfactory for muscadines. Sites where tobacco has been grown successfully are generally suitable for muscadines. Where needed to improve drainage, plant in raised rows bedded 4-12 inches high. Provisions for drainage must be made prior to planting.

Land preparation should begin the year before planting; have the soil tested, correct any deficiencies in P and K levels, and adjust the pH to 6.5 with dolomitic limestone. Use contact or systemic herbicides to clean up infestations of johnsongrass, bermudagrass, or nutsedge in the year *before* vines are set. Use a single-wire trellis, (single high wire bilateral cordon trellis), space rows 10 to 12 feet apart, and set posts 20 ft apart within rows. Plant one vine beside each post (rather than halfway between posts). This positioning will help protect the vine and minimize fruit loss during harvest, especially if mechanical harvest is planned.

Plant in late April or early May (NC and GA/SC Piedmont) after danger of spring freezes has passed, or during the winter in South GA. Commercial “grow tubes” (24-36 inches long and 3-4 inches in diameter) will accelerate growth and protect vines in the first growing season, but grow tubes may delay the development of vine cold hardiness in the fall, and for this reason they should be removed at the end of each growing season, in late September/early October. Planting depth is critical. Plant at the same depth as in the pot or nursery, in a wide, shallow hole slightly deeper than the depth of the pot. Avoid using an auger to dig planting holes, as augered holes tend to be too deep and the soil will settle (subside) over time, resulting in a vine that is planted too deep.

Pruning (training new vines)

Trellising young vines -- During the first two or three years, vines need regular training once a week throughout the growing season. Training establishes the optimal shape and position of the vine. Proper training is accomplished by selecting a single shoot and training it up a bamboo stake or string. Remove any side shoots that appear in the leaf axils by pinching them out as soon as possible, leaving the leaf intact. The objective is to have a single growing point rather than multiple shoots. Plastic vineyard tape is used for attaching vines to the bamboo stake and later to the trellis wire. Train the young vines up the stake or string to the wire, then pinch the tip back to approximately 10-12 inches below the wire. This will encourage a V-shaped branching. Choose two of these lateral branches and begin training them in opposite directions along the wire, as future “main arms” or cordons. Continually remove lateral suckers along the trunk in order to direct all the plant’s growth into the growing tips of the cordons. Until the cordons reach full length, cut back side shoots along the cordon to approximately 4 to 6 inches during the growing season. Likewise, remove all fruit at the earliest possible stage on young vines, in order to direct growth to the desired shoots.

Dormant

Winter pruning – Proper pruning aids in disease management. Hedge or hand prune vines, leaving only 3-4 inch stubs (spurs) of 1-yr-old wood protruding from the main cordon or previous season’s wood. Selectively thin these spurs or spur clusters to space them approximately 4-6 inches along the cordon. As the vines age, remove some of the older spur clusters when crowding occurs. Young vines (1-4 yrs old) require special attention to remove grape tendrils that wrap around the new cordon. If not removed, these tendrils will girdle and kill the newly-trained arm. After the 3rd or 4th year, the cordon becomes too thick for tendrils to wrap around it, and this girdling ceases to be a threat. Late winter is the best time to prune muscadines in areas subject to winter injury. **Mechanical pruning** -- Many growers use hand-held or tractor-mounted sickle bar hedge trimmers for the first rough pass in winter pruning, followed by hand pruning.

Dieback “dead arm” diseases – Fungal dieback diseases (caused by *Botryosphaeria*, *Phomopsis*, *Fusicoccum* and *Eutypa*) can quickly kill even mature plants. These diseases are often observed as plants come out of dormancy or following initial fruit swell. Symptoms are rapid death from the growing cordon tips back towards the main trunk. These fungi reside in pruning or other wounds and are present on all muscadines. Plant stress triggers disease. Any actions to reduce plant stress throughout the year will be helpful (irrigation, proper fertilization). Cold injury can predispose vines to these “dead arm” diseases by causing large, slow-healing wounds to trunks and cordons. Georgia has a state 24(c) label for the use of Topsin M to control Eutypa dead arm.

Effectiveness Ratings. The efficacy or importance of a management option is indicated by E = excellent, VG = very good, G = good, F = fair, P = poor, NA= not applicable, and UN = unknown or no data. These ratings are benchmarks; actual performance will vary.

| Pest/Problem | Management Options | Amount of Formulation per acre | Effect-iveness | Mode of Action | REI | PHI | Comments and Precautions |
|--|---|--------------------------------|----------------|----------------|--------|---------|--|
| Fruit rots and foliage diseases | Sanitation and pruning, canopy management | | G | | | | Prune to remove dead wood and old fruit pedicels that can harbor diseases. Control weeds under vines with chemicals or mowing; trim vines if needed, so shoots terminate at least 18-24 inches from the ground. This will promote air movement and drying under vines. |
| Dieback "dead arm" diseases (Botryosphaeria and other similar organisms) | Sanitation and pruning, canopy management | | G | | | | Prune to remove dead wood. Monitor vines for injury and prune to remove/replace cold-damaged or insect-damaged cordons. |
| | myclobutanil (Rally 40W) | 5 oz | UN | FRAC 3 | 24 hrs | 14 days | May be applied in 50 gallons of water per acre to protect pruning wounds. This treatment has not been evaluated on muscadines but has likely shown benefit on other grape species. Apply within 24 hrs of pruning. Do not apply more than 1.5 lb of Rally 40W per acre per year. Interval for repeat applications should not exceed 14 days. |

Prebloom (Bud break till bloom)

Angular leafspot is a fungal disease caused by the pathogen *Mycosphaerella angulata*. Leaf spotting leads to premature defoliation. When a muscadine vine defoliates prior to harvest, development of fruit ceases and grapes may fail to ripen. Total crop loss may occur. Fortunately, angular leafspot is easy to control with fungicides. In wet seasons, however, control may be expensive due to the need for repeated fungicide applications.

Black rot is a fungal disease caused by the pathogen *Guignardia bidwellii* f. sp. *Muscadinii*. Black rot can cause numerous necrotic round (ca, 10 mm) leaf spots early in the season beginning in May-June, and also causes scabby black lesions on fruit that do not penetrate the skin. Fungicides applied prior to, during and after bloom are effective in controlling black rot.

Aphids in muscadines are typically cool-season, spring pests. Natural enemies often moderate aphid numbers as the weather warms. Treat only if aphids are abundant and new shoot tips or foliage are becoming malformed.

Grape leafhoppers are sucking pests that may be very abundant, causing colorless, cleared stippling on leaves; heavy infestations can cause defoliation. Leafhoppers may be present through much of the growing season, but often in numbers that appear to do no harm. Provisional treatment thresholds are 10 leafhopper nymphs or adults per leaf based on samples from 10 vines per acre, or when leafhoppers and injury are seen (stippling and weak, unthrifty growth).

Flea beetles and May or June beetles are occasional bud and foliage feeders. The primary concern is Spring bud and shoot damage. Treat if beetles are abundant and injury is evident. Suggested treatment threshold is 5% damaged buds in a sample of 10 vines per acre. Less than 10% foliar injury is unlikely to be harmful. Beetles often feed at dusk or at night and may not be present during the day.

Prebloom (Bud break till bloom)

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|--|--|--------------------------------|---------------|----------------|--------|---------|--|
| Black rot Bitter rot Angular leaf spot Powdery mildew Macrophoma rot | myclobutanil (Rally 40W) | 3-5 oz | VG | FRAC 3 | 24 hrs | 14 days | Do not apply more than 1.5 lb of Rally 40W per acre per year. Interval for repeat applications should not exceed 14 days. Not specifically labeled for ALS |
| | thiophanate-methyl (Topsin M 70WSB) | 0.75-1.5 lb | G | FRAC 1 | 2 days | 7 days | Apply when foliage first develops and repeat at 14 to 21 day intervals or as needed. Do not apply more than 6 lb. product (4.2 lb ai)/acre/season. |
| <i>(continued on next page)</i> | | | | | | | |

Prebloom (Bud break till bloom) -- continued from previous page

| Pest/Problem | Management Options | Amount of Formulation per acre | Effect-iveness | Mode of Action | REI | PHI | Comments and Precautions |
|--|--|---------------------------------------|-----------------------|------------------------------------|--|------------|--|
| Black rot Bitter rot Angular leaf spot Powdery mildew Macrophoma rot <i>(continued)</i> | azoxystrobin (Abound 2.08SC) | 10-15.5 fl oz | E | FRAC 11 | 4 hrs | 14 days | Do not make more than five applications of Abound per acre per year, or more than two sequential sprays before alternating with other fungicides having a different mode of action. |
| | trifloxystrobin (Flint Extra) | 3.5-3.8 fl oz see label | E | FRAC 11 | 12 hrs | 14 days | Do not apply more than 23 oz of Flint per acre per season, or more than six times per season. Do not apply more than two applications of Flint before switching to a non-strobilurin fungicide. DO NOT APPLY to CONCORD GRAPES or crop injury may occur. |
| | pyraclostrobin + boscalid (Pristine WG) | 8-12.5 oz | E | FRAC 11+7 | 12 hrs (5 days for vine turning, tying) | 14 days | Do not make more than three applications of Pristine or related fungicides (strobilurin or carboxamide) per season. Do not make more than two sequential applications before alternating with a fungicide with a different mode of action. Do not use on CONCORD and related grapes (Worden, Fredonia, Niagara). |
| | kresoxim-methyl (Sovran 50 WG) | 3.2-4.8 oz | VG | FRAC 11 | 12 hrs | 14 days | Do not make more than four applications of Sovran per acre per year, or more than two sequential sprays before alternating with other fungicides having a different mode of action. |
| | EBDCs (ethylene-bis-dithiocarbamates) (Dithane M45 Manzate ProStick Maneb 75DF Penncozeb 75DF) EBDC + Zoxamide (Gavel) | 1.5-4 lb see label | G | FRAC M3 GAVEL FRAC 22+M3 | 24 hrs | 66 days | Do not apply more than 24 lb of Dithane M45, Manzate 200DF, Maneb 80WP or Penncozeb 75DF per acre per season. Repeat applications at 7 to 10 day intervals. |

(continued on next page)

Prebloom (Bud break till bloom) – continued from previous page

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|---|--|--------------------------------|---------------|----------------|---------|---|---|
| Black rot Bitter rot Angular leaf spot Powdery mildew Macrophoma rot (continued) | ziram (Ziram 76 DF) | 3-4 lb | G | FRAC M3 | 48 hrs | 21 days | Limited to a maximum of 28 lbs per acre per year. |
| | captan (Captan 50WP) | 4 lb | G | FRAC M4 | 48 hrs | 0 day | Repeat Captan applications at 7 to 14 day intervals. Apply using ground equipment in 20 to 200 gallons of water per acre. High volume dilute sprays are most effective. |
| | captan (Captan 4L) | 0.75-2 qt | G | | 48 hrs | 0 day | |
| | Triflumizole (Procure 480SC) | 4-8 fl oz | VG | FRAC 3 | 12 hrs | 7 days | Do not apply more than a total of 4 applications of Procure 480SC per crop per year. Do not apply more than a total of 32 fl oz (1 .0 lb ai) per crop per year. Apply in a minimum of 50 GPA. |
| | Fluopyram + tebuconazole (Luna Experience) | 6-8.6 fl oz | G-E | FRAC 3+7 | 12 hrs* | 14 days | *Re-entry interval is 5 days for cane tying and training activities. Rotate to a different MOA after two applications. |
| | Benzovendiflupyr (Aprovia) | 10.5 fl oz | VG | FRAC 7 | 12 hrs | 21 days | Do not exceed 3 applications per year. |
| | Isofetamid (Kenja 400SC) | 20-22 fl oz | G-E | FRAC 7 | 12 hrs | 14 days | Rotate to a different MOA after two applications. Do not apply a third application within 28 days of the second application. |
| | Azoxystrobin + flutriafol (Topguard EQ) | 5-8 fl oz | G-E | FRAC 3+11 | 12 hrs* | 14 days | Re-entry interval if 5 days for cane tying and girdling activities. |
| | Fluxapyroxad + pyraclostrobin (Mervion) | 4-5.5 fl oz | VG-E | FRAC 7+11 | 12 hrs | 14 days | Do not tank mix with any other product. May cause injury on certain bunch grape cultivars, see label. |
| | cyprodinil + fludioxonil (Switch 62.5WG) | 11-14 oz | G-E | FRAC 9+12 | 12 hrs | 7 days | Do not make more than two applications by air. |
| pydiflumetofen + fludioxonil (Miravis Prime) | 9.2-13.4 fl oz | VG-E | FRAC 7+12 | 12 hrs | 14 days | Apply on a 21-day interval. Do not make more than two applications at the maximum rate. | |

Prebloom (Bud break till bloom) – continued from previous page

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|---------------------|---|---|----------------------|-----------------------|--------------------------------|-----------------------------------|--|
| Powdery mildew ONLY | wettable sulfur (Microthiol, various brands, 80 to 92% S) | 3 to 10 lb | VG | FRAC M2 | 24 hrs | -- | Must be applied every 7-10 days. Dilute in 100 gal of water per acre. Corrosive to sprayers and trellis wires. OMRI listed for organic growers. Do not apply within two weeks of oil or oil-based products, or at temperatures above 90 F. May injure Concord and other <i>Vitis labrusca</i> cultivars. |
| Aphids | malathion 57EC | 1.5 pt | VG | IRAC 1B | 24 hrs (72 hrs for vine tying) | 3 days | Treat if aphids are abundant and terminals or foliage are becoming malformed. Aphids rarely reach damaging levels. |
| | imidacloprid (Admire Pro) | 7 to 14 fl oz (soil) 1 to 1.4 fl oz (foliar) | VG | IRAC 4A | 12 hrs | 30 days (soil) 0 days (foliar) | |
| | fenpropathrin (Danitol 2.4EC) | 5.33 to 10.66 fl oz | G | IRAC 3A | 24 hrs | 21 days | Restricted Use Pesticide. Limited to two applications per season. |
| | acetamiprid (Assail 30SG) | 2.5 oz | G | IRAC 4A | 12 hrs | 3 days | Do not make more than two applications per season. |
| Grape leafhoppers | Grape leafhopper injury is seen as colorless, cleared stippling on leaves; heavy infestations can cause defoliation. Provisional treatment thresholds suggest treating for 10 leafhopper nymphs or adults per leaf or when grape leafhoppers and injury are evident (weak, unthrifty growth.) | | | | | | |
| | malathion 57EC | 1.5 pt | VG | IRAC 1B | 24 hrs (72 hrs for vine tying) | 3 days | |
| | carbaryl (Sevin 80S) | 1.25 to 2.5 lb | G | IRAC 1A | 12 hrs | 7 days | |

(continued on next page)

Prebloom (Bud break till bloom) – *continued from previous page*

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|---|---|---|---------------|----------------|---|---|--|
| Grape leafhoppers <i>(continued)</i> | phosmet (Imidan 70W) | 1.33 to 2.12 lb | VG | IRAC 1B | 14 days | 14 days | |
| | acetamiprid (Assail 30 SG) | 2.5 oz | G | IRAC 4A | 12 hrs | 3 days | |
| | bifenthrin (Brigade 2 EC) | 3.2 to 6.4 oz | G | IRAC 3 | 12 hrs | 30 days | |
| | bifenthrin + imidacloprid (Brigadier) | 3.8 to 6.4 fl oz | G | IRAC 3+4A | 12 hrs | 30 days | |
| | fenpropathrin (Danitol 2.4EC) | 5.33 to 10.66 fl oz | G | IRAC 3A | 24 hrs | 21 days | Restricted Use Pesticide. Limited to two applications per season. |
| | imidacloprid (Admire Pro) | 7 to 14 fl oz (soil) 1 to 1.4 fl oz (foliar) | VG | IRAC 4A | 12 hrs | 30 days (soil) 0 days (foliar) | |
| | thiamethoxam (Platinum) (Actara) | 8 to 17 oz 1.5 to 3 oz | G | | 12 hrs | 60 days 5 days | Platinum is soil applied, Actara by foliar spray. Actara can be applied by ground or air. |
| | dinotefuran (Venom) | 5 to 6 oz soil 1 to 3 oz foliar | G | IRAC 4A | 12 hrs | 28 days 1 day | Only one soil application allowed per season. |
| Grape flea beetles <i>(continued on next page)</i> | carbaryl (Sevin 80WSP) | 1.25 to 2.5 lb | VG | IRAC 1A | 12 hrs | 7 days | Flea beetles chew holes in the foliage. Less than 10% foliar injury is unlikely to be harmful. Treat if flea beetles are abundant and injury is evident. |
| | Malathion 57EC | 1.5 pt | G | IRAC 1B | 24 hrs (72 hrs for vine tying) | 3 days | Grape flea beetle larvae also feed on buds. Treatment is recommended if 5% of vines exhibit bud damage. |
| | cyfluthrin (Baythroid 2 EC) | 2.4-3.2 fl oz | G | IRAC 3A | 12 hrs | 3 days | |

Prebloom (Bud break till bloom) – *continued from previous page*

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|--|--|---------------------------------------|----------------------|-----------------------|------------|------------|---|
| Grape flea beetles <i>(continued)</i> | phosmet (Imidan 70-W) | 1.33-2.125 lb | G | IRAC 1B | 14 days | 14 days | The 14-day REI may make this product impractical to use for most growers. |
| | bifenthrin +imidacloprid (Brigadier) | 3.8 to 6.4 fl oz | G | IRAC 3+4A | 12 | 30 | Restricted Use Pesticide. |
| | fenpropathrin (Danitol 2.4EC) | 5.33 to 10.66 fl oz | G | IRAC 3A | 24 hrs | 21 days | Restricted Use Pesticide. Limited to two applications per season. |
| Climbing cutworms | fenpropathrin (Danitol 2.4EC) | 5.33 to 10.66 fl oz | G | IRAC 3A | 24 hrs | 21 days | Restricted Use Pesticide. Limited to two applications per season. |
| | carbaryl (Sevin 80WSP) | 2.5 lbs | VG | IRAC 1A | 12 hrs | 7 days | Use of Sevin may encourage mites. Do not concentrate spray on clusters or visible residue may result. Do not apply more than 12 ½ lb/acre per year. |
| | carbaryl (Sevin XLR Plus) | 2 qt | VG | | 12 hrs | 7 days | Repeat applications as necessary up to a total of 5 times but not more often than once every 7 days. Do not apply more than 10 qt/acre per year. |
| | cyfluthrin (Baythroid 2EC) | 2.4-3.2 fl oz | G | IRAC 3A | 12 hrs | 3 days | |
| | <i>(continued on next page)</i> | methoxyfenozide (Intrepid 2F) | 12-16 fl oz | G | IRAC 18 | 4 hrs | 30 days |

Prebloom (Bud break till bloom) – continued from previous page

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|----------------------------------|--|--------------------------------|---------------|----------------|--------|---------|--|
| Climbing Cutworms (continued) | spinetoram (Delegate 25 WG) | 3-5 oz | G | IRAC 5 | 4 hrs | 7 days | |
| | flubendiamide + buprofezin (Tourismo 12.5) | 10-14 fl oz | F | IRAC 16+28 | 12 hrs | 7 days | |
| | spinosad (Entrust 80W) | 1.25-2.5 oz | G | IRAC 5 | 4 hrs | 7 days | OMRI approved. |
| | <i>Bacillus thuringiensis</i> [Bt] (Dipel DF and others) | 0.5-1 lb | F | IRAC 11 | 4 hrs | 0 days | OMRI approved. |
| | rynaxypyr (Altacor) | 2.0-4.5 oz | G | IRAC 28 | 4 hrs | 14 days | Use between 100-200 gallons per acre total spray volume. |

Bloom (Do Not Apply Insecticides During Bloom) – continued from previous page

| Pest/Problem | Management Options | Amount of Formulation per acre | Effect-iveness | Mode of Action Code | REI | PHI | Comments and Precautions |
|---|---|--------------------------------|----------------|--|--------------------------|--|--|
| <p>The disease Ripe rot (caused by <i>Colletotrichum</i> sp. fungi) is one of the most difficult fungal pathogens to control on muscadine grapes. Most bronze-fruited cultivars are susceptible, while most purple-fruited muscadines are resistant. Fungicidal control has been most successful when strobilurin-type fungicides [FRAC code 11] are combined or alternated with fungicides having a different mode of action. Sprays should begin at bloom or immediately following bloom (first cover). Ripe rot is most severe in wet harvest seasons and when fruit is left hanging too long on the vine. Timely harvest and rapid cooling of harvested fruit aids in control.</p> | | | | | | | |
| Black rot Bitter rot Angular leaf spot Powdery mildew Macrophoma rot (continued) | EBDCs (ethylene-bis-dithiocarbamates) Dithane M45 Manzate ProStick Maneb 75DF Penncozeb 75DF EBDC + Zoxamide (Gavel) | 1.5-4 lb see label | G | FRAC M3 GAVEL FRAC 22+M3 | 24 hrs 48 hrs | 66 days 21 days 0 day 0 day | Do not apply more than 24 lb Dithane M45, Manzate 200DF, Maneb 80WP or Penncozeb 75DF per acre per season. Repeat applications at 7 to 10 day intervals. Repeat Captan applications at 7 to 14 day intervals. Do not apply 50WP by air. |
| | ziram (Ziram 76 DF) | 3-4 lb | G | FRAC M3 | 48 hrs | 21 days | |
| | captan (Captan 50WP) 80 WP Various | 4 lb see label | G | FRAC M4 | 48 hrs | 0 day | |
| | captan (Captan 4L) | 0.75-2 qt | G | | 48 hrs | 0 day | |
| | Triflumizole (Procure 480SC) | 4-8 fl oz | VG | FRAC 3 | 12 hrs | 7 days | Do not apply more than a total of 4 applications of Procure 480SC per crop per year. Do not apply more than a total of 32 fl oz (1 .0 lb ai) per crop per year. Apply in a minimum of 50 GPA. |
| | Fluopyram + tebuconazole (Luna Experience) | 6-8.6 fl oz | G-E | FRAC 3+7 | 12 hrs* | 14 days | *Re-entry interval is 5 days for cane tying and training activities. Rotate to a different MOA after two applications. |
| | Benzovendiflupyr (Aprovia) | 10.5 fl oz | VG | FRAC 7 | 12 hrs | 21 days | Do not exceed 3 applications per year. |
| | (continued on next page) | Isfetamid (Kenja 400SC) | 20-22 fl oz | G-E | FRAC 7 | 12 hrs | 14 days |

Bloom (Do Not Apply Insecticides During Bloom) – continued from previous page

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action Code | REI | PHI | Comments and Precautions |
|----------------------------------|---|---------------------------------------|----------------------|--|------------|------------|---|
| Ripe rot (<i>continued</i>) | EBDCs (ethylene-bis-dithiocarbamates) Dithane M45 Manzate ProStick Maneb 75DF Penncozeb 75DF EBDC + Zoxamide (Gavel) | 1.5-4 lb see label | G | FRAC M3 GAVEL FRAC 22+M3 | 24 hrs | 66 days | Do not apply more than 24 lb Dithane M45, Manzate 200DF, Maneb 80WP or Penncozeb 75DF per acre per season. Repeat applications at 7 to 10 day intervals. |
| | ziram (Ziram 76 DF) | 3-4 lb | G | FRAC M3 | 48 hrs | 21 days | |
| | captan (Captan 50WP) 80 WP Various | 4 lb see label | G | FRAC M4 | 48 hrs | 0 day | Repeat Captan applications at 7 to 14 day intervals. Do not apply 50WP by air. |
| | captan (Captan 4L) | 0.75 to 2 qt | G | | 48 hrs | 0 day* | |
| Powdery mildew ONLY | wettable sulfur (Microthiol, various brands, 80 to 92% S) | 3-10 lb | VG | FRAC M2 | 24 hrs | -- | Must be applied every 7-10 days. Dilute sulfur in 100 gal of water per acre. Sulfur is corrosive to sprayers and trellis wires. Do not apply within two weeks of oil or oil-based products, or at temperatures above 90 F. May injure Concord and other <i>Vitis labrusca</i> cultivars. Some but not all sulfur products are OMRI listed for use in organic production. Check the label! |

First Cover (post-bloom) – continued from previous page

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|--|--|---|---------------|------------------------------------|-----------|---------|--|
| Black rot Bitter rot Angular leaf spot Powdery mildew Macrophoma rot <i>(continued)</i> | kresoxim-methyl (Sovran 50 WG) | 3.2 to 4.8 oz | VG | FRAC 11 | 12 hrs | 14 days | Do not make more than four applications of Sovran per acre per year. Do not apply more than two sequential sprays of Sovran. Alternate applications with other fungicides which have a different mode of action. |
| | EBDCs (ethylene-bis-dithiocarbamates) Mancozeb, Dithane M45 Manzate ProStick Maneb 75DF Penncozeb 75DF EBDC + Zoxamide (Gavel) | 1.5 to 4 lb | G | FRAC M3 GAVEL FRAC 22+M3 | 24 hrs | 66 days | Do not apply more than 24 lb of Dithane M45, Manzate 200DF, Maneb 80WP or Penncozeb 75DF per acre per season. Repeat applications at 7 to 10 day intervals. **Note PHI of 66 days – cannot be used post-bloom for early-ripening cultivars. |
| | Ziram 76DF | 3 to 4 lb | G | FRAC M3 | 48 hrs | 21 days | |
| | captan (Captan 50WP) | 4 lb | G | FRAC M4 | 48 hrs | 0 day* | Repeat Captan applications at 7 to 14 day intervals. Label allows application up to day of harvest; however; note 48 hr re-entry interval. |
| | captan (Captan 4L) | 0.75 to 2 qt | G | | 48 hrs | 0 day* | |
| | Triflumizole (Procure 480SC) | 4-8 fl oz | VG | FRAC 3 | 12 hrs | 7 days | Do not apply more than a total of 4 applications of Procure 480SC per crop per year. Do not apply more than a total of 32 fl oz (1 .0 lb ai) per crop per year. Apply in a minimum of 50 GPA. |
| | Fluopyram + tebuconazole (Luna Experience) | 6-8.6 fl oz | G-E | FRAC 3+7 | 12 hrs* | 14 days | *Re-entry interval is 5 days for cane tying and training activities. Rotate to a different MOA after two applications. |
| | Benzovendiflupyr (Aprovia) | 10.5 fl oz | VG | FRAC 7 | 12 hrs | 21 days | Do not exceed 3 applications per year. |
| | Isofetamid (Kenja 400SC) | 20-22 fl oz | G-E | FRAC 7 | 12 hrs | 14 days | Rotate to a different MOA after two applications. Do not apply a third application within 28 days of the second application. |
| | <i>(continued on next page)</i> | Azoxystrobin + flutriafol (Topguard EQ) | 5-8 fl oz | G-E | FRAC 3+11 | 12 hrs* | 14 days |

First Cover (post-bloom) – continued from previous page

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action Code | REI | PHI | Comments and Precautions |
|---|--|---------------------------------|---------------|---------------------|--|---------|--|
| Ripe rot (continued) | ziram (Ziram 76 DF) | 3-4 lb | G | FRAC M3 | 48 hrs | 21 days | |
| | captan (Captan 50WP) 80 WP Various | 4 lb see label | G | FRAC M4 | 48 hrs | 0 day | Repeat Captan applications at 7 to 14 day intervals. Do not apply 50WP by air. |
| | captan (Captan 4L) | 0.75 to 2 qt | G | | 48 hrs | 0 day* | |
| Powdery mildew ONLY (Sulfur will not control other diseases) | wettable sulfur (various brands, 80 to 92% S) | 2 to 5 lbs | VG | FRAC M2 | 24 hrs | -- | Must be applied every 7-10 days. Dilute sulfur in 100 gal of water per acre. Sulfur is corrosive to sprayers and trellis wires. Do not apply within two weeks of oil or oil-based products, or at temperatures above 90 F. May injure Concord and other <i>Vitis labrusca</i> cultivars. |
| Sooty blotch (Primarily seen on the cultivar 'Fry') | captan (Captan 50WP) (Captan 4L) | 2.0 to 4.0 lb 0.75 to 2.0 qt | G | FRAC M4 | 48 hrs | 0 days | *Label allows application up to day of harvest; however; note 48 hr re-entry interval. |
| | azoxystrobin (Abound 2.08SC) | 10.0 to 15.5 fl oz | E | FRAC 11 | 4 hrs | 14 days | Do not make more than five applications of Abound per acre per year. Do not apply more than two sequential sprays of Abound. Alternate applications with other fungicides which have a different mode of action. |
| | trifloxystrobin (Flint Extra) | 3.5 to 3.8 oz | E | FRAC 11 | 12 hrs | 14 days | Do not apply more than 23 oz of Flint Extra per acre per season. Do not make more than six applications of Flint per season. Do not apply more than two applications of Flint before switching to a non-strobilurin fungicide. |
| | pyraclostrobin + boscalid (Pristine) | 8.0 to 12.5 oz | E | FRAC 11+7 | 12 hrs (5 days for vine turning, tying) | 14 days | Do not make more than five applications of Pristine or related fungicides (FRAC 7 or 11) per season. Do not make more than two sequential applications of Pristine before alternating with a fungicide with a different mode of action. |

Summer cover (post-bloom) sprays until harvest

Fruit rots are best controlled by fungicide applications applied from pre-bloom to first cover. Proper pruning and removal of infected berries, leaves and old fruit stems from the previous winter may reduce disease. If **bitter rot** is very severe, fungicides can be applied at shoot emergence, 3-5 inches, 8-10 inches and at 7-10 day intervals until fruit set. **Black rot**-susceptible cultivars can be sprayed with a fungicide every 14 days from the start of new growth until after bloom. **Ripe rot** can be severe on susceptible cultivars such as Higgins, Magnolia, Summit, Watergate, Carlos, Fry, Dixieland, and Scuppernong. Abound and other QoI fungicides are the materials of choice in vineyards where ripe rot is a problem. However, ripe rot resistance is likely developing to these chemicals. **Sooty blotch** causes a dark, superficial discoloration on the surface of otherwise healthy fruit; the disease is common on unsprayed vines of the cultivar Fry. **Macrophoma rot** control can be improved with early cover sprays of Captan, particularly on susceptible cultivars such as Cowart, Fry, Higgins, Summit and Triumph.

Cover spray insecticides are often not needed in muscadines due to moderate insect pressure. Growers should rely on “as-needed” applications, while using the IPM approach of scouting frequently for insect damage, including detailed plant examination of 10 vines per acre for insect or mite pests.

Grape berry moth is somewhat sporadic, but if present can be very damaging. This moth has several generations per season, beginning around bloom. In muscadines, grape berry moth is normally a mid- to late-season pest. Eggs are laid on the berry clusters, and young larvae enter berries at the stem end; their feeding, webbing and frass can damage multiple berries within a cluster. Spray promptly if infested clusters are seen or if a vineyard has a history of grape berry moth infestations.

Grape curculio is a small weevil (snout beetle) that typically emerges in mid-June (in GA). Grape curculio initially feeds on the underside of leaves in a shallow zigzag pattern before beginning to lay eggs in the berries. Spray if grape curculio are present or if either foliar or fruit injury are seen. This is a sporadic pest that is sometimes a problem in unmanaged vineyards where weeds and wild hosts thrive.

Green June beetle and **Japanese beetle** populations can get out of hand rapidly. Moderate defoliation by Japanese beetles is seldom damaging, but in some years these insects can be found feeding on flowers, and this can greatly reduce the crop. Fruit feeding by green June beetle is serious, so do not allow these pests to feed heavily on and become abundant in blocks with ripe fruit. Beware of heavy emergence and migration to blocks with ripe fruit after rains. Multiple applications are often necessary to maintain control if populations are allowed to build up. Ripening fruit and aggregation pheromone may quickly attract ‘new’ beetles to your vineyard after successful applications.

Wasps, hornets, yellowjackets and imported fire ants may be attracted to ripening fruit. Feeding injury or the presence of insect debris in harvested fruit can be damaging, and stinging insects discourage pick-your-own customers. Pre-harvest insecticide options are limited; monitor and treat only as needed.

Spider mites and others – Mites are capable of explosive population growth. Treat if more than 10 mites per leaf are found or if mites are present and leaves are webbed or bronzed. Drought and heavy crop load aggravate mite injury, especially early in a growing season. Two treatments at one week intervals may be needed if using a material that only kills adult mites. **Brevipalpid** mites (flat mites) cause russetting around leaf veins and at the stem end of berries in NC, and this mite may occasionally warrant control.

Stink bugs – The impact of stink bugs and other piercing/sucking insects in muscadine grapes is unclear, but their feeding may cause fruit to shrivel, spot or abort. No threshold has been developed for these pests.

Summer cover (post-bloom) sprays until harvest— *continued from previous page*

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|--|---|--------------------------------|---------------|----------------|---------|---------|--|
| Black rot Bitter rot Ripe rot Macrophoma rot Angular leaf spot Sooty blotch | Same as sprays for First Cover , except EBDCs must now be excluded due to their 66 day preharvest interval | | | | | | Exclude EBDCs (Maneb, Manex, Penncozeb, Dithane M-45, Gavel and similar). Other effective cover sprays should include tank mixes of Topsin M or Rally with Captan or Captec, OR products containing two active ingredients (two modes of action) |
| Stink bugs | phosmet (Imidan 70W) | 1.33 to 2.125 lb | VG | IRAC 1B | 14 days | 14 days | Imidan seldom prompts outbreaks of secondary pests such as mites. The 14-day REI may make this product impractical to use for most growers. |
| | fenpropathrin (Danitol 2.4EC) | 10 to 21 fl oz | VG | IRAC 3A | 24 hrs | 21 days | Danitol is a pyrethroid, its use may prompt rebound of mite numbers later in the season. |
| Japanese beetle | carbaryl (Sevin 80WSP) | 1.25 to 2.5 lbs | VG | IRAC 1A | 12 hrs | 7 days | Use of Sevin may encourage mites. Do not concentrate spray on clusters or visible residue may result. Do not apply more than 12 ½ lb/acre per year. |
| | carbaryl (Sevin XLR Plus) | 1 to 2 qt | VG | | 12 hrs | 7 days | Repeat applications as necessary up to a total of 5 times but not more often than once every 7 days. Do not apply more than 10 qt/acre per year. |
| | fenpropathrin (Danitol 2.4EC) | 10 to 21 fl oz | E | IRAC 3A | 24 hrs | 21 days | Danitol is a pyrethroid, its use may prompt rebound of mite numbers later in the season. |
| | acetamiprid (Assail 30SG) | 2.5 oz | G | IRAC 4A | 12 hrs | 3 days | Do not make more than two applications per season. |
| | imidacloprid (Admire Pro) | 1.3 to 1.4 fl oz | VG | | 12 hrs | 0 days | Do not apply by air. |
| | clothianidin (Clutch 50WDG) | 3 oz | G | | 12 hrs | 0 days | Do not make more than two applications per season. |

(continued on next page)

Summer cover (post-bloom) sprays until harvest – *continued from previous page*

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|---------------------------------------|---|--------------------------------|---------------|----------------|--------|--------------|--|
| Japanese beetle <i>(continued)</i> | phosmet (Imidan 70 WP) | 1.33 to 2.125 lb | G | IRAC 1B | 14 hrs | 7 or 14 days | Seven-day PHI at the lowest rate, 14-day PHI for rates above 1.33 lb. Do not apply more than 6.5 lbs per acre per year. |
| | Surround | 25 to 50 lbs | G | none | 4 hrs | 0 days | Surround coats leaves and fruit, rendering them unattractive to Japanese beetles. Surround may also decrease other pest activity. Surround is OMRI listed for use by organic growers. Visible residue may persist on fruit at harvest. |
| Grape berry moth | carbaryl (Sevin 80WSP) | 2.5 lbs | VG | IRAC 1A | 12 hrs | 7 days | Use of Sevin may encourage mites. Do not concentrate spray on clusters or visible residue may result. Do not apply more than 12 ½ lb/acre per year. |
| | carbaryl (Sevin XLR Plus) | 2 qt | VG | | 12 hrs | 7 days | Repeat applications as necessary up to a total of 5 times but not more often than once every 7 days. Do not apply more than 10 qt/acre per year. |
| | clothianidin (Clutch 50 WDG) | 3 oz | G | IRAC 4 | 12 hrs | 0 days | |
| | spinosad (Entrust 80W) (Entrust 2SC) | 1.25 to 2.5 oz 4 to 8 fl oz | G | IRAC 5 | 4 hrs | 7 days | Treatments should be timed to egg hatch. Entrust is OMRI listed. |
| | spinetoram (Delegate) | 3-5 oz | G | | 4 hrs | 3 days | |
| | methoxyfenozide (Intrepid 2F) (Intrepid Edge) | 8 to 16 fl oz 6 to 12 fl oz | G | IRAC 18 | 4 hrs | 30 days* | Treatments should be timed to egg hatch. *Shorter preharvest intervals are permitted when reduced rates are used, see label. |
| | bifenthrin (Brigade 2 EC) | 3.2 to 6.4 fl oz | G | IRAC 3 | 12 hrs | 30 days | |
| | bifenthrin + imidacloprid (Brigadier) | 3.8 to 6.4 fl oz | G | IRAC 3+4A | 12 hrs | 30 days | |
| | indoxacarb (Avaunt) | 5 to 6 oz | G | IRAC 22 | 12 hrs | 7 days | |

(continued on next page)

Summer cover (post-bloom) sprays until harvest – *continued from previous page*

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|---|---|--------------------------------|---------------|----------------|--------|--------------|---|
| Grape berry moth (<i>continued</i>) | phosmet (Imidan 70 WP) | 1.33 to 2.125 lb | G | IRAC 1B | 14 hrs | 7 or 14 days | Seven-day PHI at the lowest rate, 14 day PHI for rates above 1.33 lb. Do not apply more than 6.5 lbs per acre per year. |
| | methomyl (Lannate SP) | 0.5 to 1 lb | G | IRAC 1A | 7 days | 1 or 14 days | Preharvest interval is 1 day for fresh market grapes and 14 days for wine grapes. |
| | fenpropathrin (Danitol 2.4 EC) | 10.66 to 21.33 fl oz | G | IRAC 3 | 24 hrs | 21 days | Do not exceed 2.66 pints per acre per season. Make no more than two applications per season. |
| | pyriproxyfen (Esteem 0.83 EC) | 16 fl oz | G | IRAC 7C | 12 hrs | 21 days | |
| | Rynaxypyr (Altacor) | 3 to 4.5 fl oz | G | IRAC 28 | 4 hrs | 14 days | Use between 100 and 200 gallons per acre total spray volume. |
| Grape rootworm, Southern grape rootworm | Grape rootworm larvae feed on roots. Adults are small, black weevils that make distinctive chain-like feeding markings on leaves. Foliar feeding does not result in yield reduction, but root feeding may reduce plant vigor over time. Treatments should be timed to adult activity, which typically peaks in June or July. Grape rootworms are sporadic pests and should not be treated preventatively. | | | | | | |
| | carbaryl (Sevin 80WSP) | 2.5 lbs | VG | IRAC 1A | 12 hrs | 7 days | Use of Sevin may encourage mites. Do not concentrate spray on clusters or visible residue may result. Do not apply more than 12 ½ lb/acre per year. |
| | carbaryl (Sevin XLR Plus) | 2 qt | VG | | 12 hrs | 7 days | Repeat applications as necessary up to a total of 5 times but not more often than once every 7 days. Do not apply more than 10 qt/acre per year. |
| Scale insects | Scale insects are occasional pests of muscadine grapes. | | | | | | |
| | buprofezin (Applaud 70 DF) | 9 to 12 oz | G | IRAC 16 | 12 hrs | 7 days | Apply when crawlers are observed. |
| | spirotetramat (Movento 2 SC) | 6 to 8 fl oz | G | IRAC 23 | 24 hrs | 7 days | |

Summer cover (post-bloom) sprays until harvest – *continued from previous page*

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|--------------|--|------------------------------------|---------------|----------------|------------------|---------|---|
| Spider mites | Sample for mites using a minimum 10x hand lens. There is no clearly defined threshold for mites in muscadine grapes. Treatment for vinifera grapes is recommended when greater than 50% of the leaves are infested. Fast-moving predatory mites can be distinguished from slower spider mites. Some insecticides such as carbaryl (Sevin), bifenthrin (Brigade), and fenpropathrin (Danitol) can flare mite populations. Use with caution when mites are present. Rotate acaricides (miticides) between modes of action (MOAs) to minimize selection for resistance. | | | | | | |
| | pyridaben (Nexter 75WP) | 4.4 oz | VG | IRAC 21 | 12 hrs | 7 days | Nexter is an excellent miticide for European red mite. |
| | bifenazate (Acramite 50WS) | 0.75-1 lb | VG | IRAC UN | 12 hrs to 5 days | 14 days | Acramite kills eggs, larvae and adult spider mites. Allow at least a week before considering a second application based on scouting. |
| | abamectin (Abba 0.15 EC, Agri-Mek 0.15EC and others) Agri-Mek SC | 8 – 16 fl oz 1.75-3.5 fl oz | E | IRAC 6 | 12 hrs to 4 days | 28 days | Apply with a non-ionic surfactant. Test for possible cultivar-specific phytotoxicity prior to spraying entire blocks. Do not reapply within 21 days. Do not apply by air. Do not make more than two applications per season. This adulticide, will not impact eggs or larvae. Restricted Use Pesticide. |
| | fenpyroximate (Portal XLO, Fujimite 5 EC) | 2 pt | E | IRAC 21 | 12 hrs | 14 days | Portal and Fujimite kill eggs, larvae and adult spider mites. Allow at least a week before considering a second application based on scouting. Do not make more than two applications per season. Do not apply by air. |
| | etoxazole (Zeal) | 3 oz | G | IRAC 10B | 12 hrs | 28 days | Zeal is a growth regulator that kills eggs and young mites. It is most effective if applied when mite populations are low. |
| | cyflumetofen (Nealta) | 13.7 fl oz | VG | IRAC 25 | 12 hrs | 14 days | Do not make more than two Nealta applications per season. Rotate to another mode of action between applications. |
| | <i>(continued on next page)</i> | | | | | | |

Summer cover (post-bloom) sprays until harvest – *continued from previous page*

| Pest/Problem | Management Options | Amount of Formulation per acre | Effect-iveness | Mode of Action | REI | PHI | Comments and Precautions |
|------------------------------------|--|--------------------------------|----------------|----------------|--------|---------|--|
| Spider mites <i>(continued)</i> | Horticultural oils (many materials, including Glacial Spray Fluid) | 1 to 2% by volume | G | UNK | 4 hrs | 0 days | Some oils are OMRI listed. Do not use in combination or immediately before/after Captan or sulfur-containing products. Do not use with carbaryl or dimethoate. Do not use with any product whose label recommends the use of no oils. Do not use in combination with foliar fertilizers. |
| | spirodiclofen Envidor 2 SC | 18 fl oz | G | IRAC 23 | 12 hrs | 14 days | |
| | fenbutatin-oxide (Vendex 50WP) | 1 lb | F | IRAC 12B | 48 hrs | 28 days | Do not retreat with Vendex for 21 days. Restricted Use Pesticide. |
| Grape tumid gallmaker | Grape tumid gallmaker adults are small flies. Their larvae, or maggots, infest clusters, leaves, or stems and cause the plant to create galls. Galls can be quite large and are often reddish. Infestations are usually localized and do not typically require chemical treatment. | | | | | | |
| | Hand removal | NA | G | NA | NA | NA | Removal of affected plant parts is generally sufficient to prevent further infestation. |
| | spirotetramat (Movento 2 SC) | 6 to 8 fl oz | G | IRAC 23 | 24 hrs | 7 days | |

Grape Root Borer Control

| Pest/Problem | Management Options | Amount of Formulation per acre | Effectiveness | Mode of Action | REI | PHI | Comments and Precautions |
|------------------|---|--------------------------------|---------------|-------------------|-----|-----|---|
| Grape root borer | Grape root borer is potentially the most significant pest of grapes in North Carolina, but they are not necessarily present in all vineyards. Grape root borer moths should be monitored with pheromone baited traps. Trap captures may begin in July and continue through September. If moths are confirmed within a vineyard, mating disruption is the most effective control tool. | | | | | | |
| | Mounding | | G | | | | Mounding is an organically acceptable practice. Mounds need to be knocked down before winter. |
| | Isomate GRB | 100 Dispensers | VG | Mating disruption | | | Pheromone-based mating disruption has provided 90% reduction of pupal counts on bunch grapes in Virginia. Growers in NC have reported near 100% disruption (based on reduced trap catches of adult males) in muscadine vineyards. |

Fire ant control

Fire ants – Fire ants may be controlled through the use of bait products. These products are often not fast-acting but operate by sterilizing the colony (the queen lays sterile eggs). They must be applied when the ants are actively foraging -- test for foraging behavior by placing food (hot dog pieces, potato chips, sardines, etc.) and checking 30 min later for ant activity. Apply during dry weather; moisture reduces the effectiveness of bait products. Insecticides applied for control of other insect pests will also aid in suppression of fire ants. Fire ant populations are at their highest in young vineyards, but the number of mounds drops rapidly after the vines develop a mature canopy that shades the vineyard floor

| | | | | | | | |
|-----------------------|--|----------------|----|---------|--------|-------|--|
| Red Imported Fire Ant | pyriproxyfen (Esteem Ant Bait) | 1.5-2.0 lbs | VG | IRAC 7C | 12 hrs | 1 day | For grapes do not exceed 0.22 lbs active ingredient (ai, pyriproxyfen) per season. |
| | methoprene (Extinguish Professional Fire Ant Bait) | 1.0 to 1.5 lbs | VG | IRAC 7A | 4 hrs | 0 day | Labeled for use on cropland, including grape vineyards. |

Efficacy of selected fungicides against diseases of muscadine grape (E = excellent, VG = very good, G = good, F = fair, P = poor, NA = not recommended, UN = control unknown) These ratings are benchmarks, actual performance will vary.

| Fungicide | PHI (days) | FRAC MOA | Bitter rot | Powdery mildew | Ripe rot | Macro-phoma rot | Black rot | Sooty blotch | Dead arm | Angular leaf spot |
|---|-----------------------------|----------|------------|----------------|----------|-----------------|-----------|--------------|----------|-------------------|
| Thiophanate-methyl (Topsin- M) | 7 days | 1 | G | G | F | G | G | G | F | G |
| Myclobutanil (Rally) | 14 days | 3 | G | VG | NA | G | VG | G | UN | VG |
| Triflumizole (Procure) | 7 days | 3 | VG | VG | UN | VG | VG | UN | UN | VG |
| Fluopyram + tebuconazole (Luna Experience) | 14 days | 3+7 | VG | VG | UN | G | VG | UN | UN | E |
| Benzovendiflupyr (Aprovia) | 21 days | 7 | VG | UN | G | VG | VG | UN | UN | VG |
| Isofetamid (Kenja) | 14 days | 7 | VG | VG | G | E | VG | UN | UN | VG |
| Azoxystrobin + flutriafol (Topguard EQ) | 14 days | 3+11 | E | VG | G | VG | VG | UN | UN | E |
| Fluxapyroxad + pyraclostrobin (Mervion) | 14 days | 7+11 | E | VG | VG | VG | VG | UN | UN | E |
| Pyraclostrobin + boscalid (Pristine) | 14 days | 7+11 | G | VG | VG | E | VG | E | F | VG |
| Kresoxim-methyl (Sovran) | 14 days | 11 | G | G | G | G | G | G | F | G |
| Azoxystrobin (Abound) | 14 days | 11 | G | VG | VG | VG | VG | VG | F | VG |
| Trifloxystrobin (Flint Extra) | 14 days | 11 | G | VG | VG | E | VG | E | F | G |
| cyprodinil + fludioxonil (Switch) | 7 days | 9+12 | E | VG | VG | E | VG | UN | UN | G |
| pydiflumetofen + fludioxonil (Miravis Prime) | 14 days | 7+12 | E | VG | VG | E | VG | UN | UN | VG |
| Copper oxychloride + Copper hydroxide (Badge) | 0 day (48 hrs re-entry) | M 1 | NA | G | NA | NA | G | UN | UN | VG |
| Wettable Sulfur (Microthiol and other trade names) | 0 day (24 hrs re-entry) | M 2 | NA | VG | NA | NA | NA | F | NA | NA |
| Mancozeb + Zoxamide (Gavel) | 66 days | 22+M3 | G | G | NA | G | G | UN | UN | G |
| Ziram (Ziram) | 21 days | M 3 | G | G | G | G | G | G | F | G |
| Captan (Captan, Captec) | 0 days (72 hrs re-entry) | M 4 | G | G | VG | G | G | G | F | G |
| EBDCs (includes Maneb, Manex, Penncozeb, Manzate, Dithane M-45) | 66 days | M 3 | G | G | G | G | G | G | F | G |

Seasonal ‘at a glance’ fungicidal spray schedule -- options for muscadine grape

Do not rely on a single product -- use alternating modes of action (MOA) shown below in brackets [FRAC Code]. EARLY SPRAYS are more effective than late-season sprays. This table is only a guide – read and follow the label prior to making applications.

| Developmental Stage  | Dormant | Pre-bloom -- new shoots to 6 inches | Second pre-bloom | Bloom – 50% open flowers | First Cover – small green fruit | Subsequent Cover sprays (second, third, fourth) | Can be used up to 14 days preharvest | Can be used up to 7 days preharvest |
|---|---|--|--|--|---|---|---|-------------------------------------|
| Diseases (bold) Fungicide [FRAC Code] | Dieback "dead arm" Eutypa, Botryosphaeria and similar organisms | These products can be used to manage most fruit rots and leaf diseases (Black rot, Bitter rot, Angular leaf spot, Powdery mildew, Macrophoma rot) but do NOT provide control of Ripe rot (<i>Colletotrichum</i> sp.) | | | | | | |
| | | Topsin M [1] Rally [3] Procure [3] Luna Experience [3+7] | Topsin M [1] Rally [3] Procure [3] Luna Experience [3+7] | Topsin M [1] Rally [3] Procure [3] Luna Experience [3+7] | Topsin M [1] Rally [3] Procure [3] Luna Experience [3+7] | Topsin M [1] Rally [3] Procure [3] Luna Experience [3+7] | Topsin M [1] Rally [3] Procure [3] Luna Experience [3+7] | Topsin M [1] Procure [3] |
| | Rally [3] applied at pruning time to fresh cuts | These products can be used to manage fruit rots and leaf diseases (Black rot, Bitter rot, Angular leaf spot, Powdery mildew, Macrophoma rot) – INCLUDING management of Ripe rot (<i>Colletotrichum</i> sp.) | | | | | | |
| | | Aprovia [7] Kenja [7] Topguard EQ [3+11] Mervion [7+11] Pristine [7+11] Sovran [11] Abound [11] Flint Extra [11] Switch [9+12] Miravis Prime [7+12] Ziram [M3] Captan [M4] EBDCs (66-day preharvest interval) -- includes Maneb, Manex, Penncoze, Manzate, Dithane M-45 [M3] and Gavel [22+M3] | Aprovia [7] Kenja [7] Topguard EQ [3+11] Mervion [7+11] Pristine [7+11] Sovran [11] Abound [11] Flint Extra [11] Switch [9+12] Miravis Prime [7+12] Ziram [M3] Captan [M4] EBDCs (66-day preharvest interval) -- includes Maneb, Manex, Penncoze, Manzate, Dithane M-45 [M3] and Gavel [22+M3] | Aprovia [7] Kenja [7] Topguard EQ [3+11] Mervion [7+11] Pristine [7+11] Sovran [11] Abound [11] Flint Extra [11] Switch [9+12] Miravis Prime [7+12] Ziram [M3] Captan [M4] (Note -- Do not use EBDCs within 66 days of harvest) | Kenja [7] Topguard EQ [3+11] Mervion [7+11] Pristine [7+11] Sovran [11] Abound [11] Flint Extra [11] Switch [9+12] Miravis Prime [7+12] Captan [M4] (Note – fungicides will leave a visible residue on harvested fruit when used close to harvest) | Switch [9+12] Captan [M4] | | |
| | | Powdery mildew ONLY | | | | | | |
| | | Wettable Sulfur [M2] | Wettable Sulfur [M2] | Wettable Sulfur [M2] | | | | |

Weed Management Grape Vineyards

The primary goal of any weed management program is to minimize competition in order to direct as many resources, like water, nutrients, and light, as possible toward crop growth. It is essential to minimize or eliminate competition in newly planted and young vineyards so that vine growth can be maximized to bring that vineyard into productivity as soon as possible. Research has shown that failure to control weeds through July in newly planted vineyards will reduce vine growth and may increase vine mortality due to water stress. In older, established vineyards competition will reduce grape yields. The weed management programs outlined in this publication are designed to control weeds at levels to prevent competition and maximize fruit yields.

Herbicide Resistance Management

The development of herbicide resistant weed species has increased significantly across the Southeast during the past few years. Lately weed resistance to glyphosate has been the most common resistance development which is largely related to the widespread planting of glyphosate resistant crops. The utilization of herbicides with differing modes of action (MOA) during the growing season or tank mixing herbicides with differing MOA are strategies that can be utilized to prevent the development of herbicide resistant weeds. As a means to assist growers with identifying herbicides having like MOA a number system identifying herbicides by MOA has been developed and is being utilized. In the table below there is a MOA number for each herbicide active ingredient to aid growers in making management decisions that will prevent the development of herbicide resistance or address options for managing a known resistant weed population that may be in or near the vineyard.

Additionally, growers are encouraged to find at least two herbicide programs containing different herbicides to rotate on an annual basis. By rotating herbicide programs growers not only minimize the risk of herbicide resistance developing but they also minimize the likelihood of selecting for weeds that one herbicide program may not be particularly effective at controlling.

Vineyard Herbicide Options

| Weed/Timing | Material | Amount of Formulation per Acre | Crop Age Restrictions | REI (hrs) | Comments |
|---|---|--------------------------------|--|-----------|---|
| PREPLANT/ SITE PREPARATION | | | | | |
| PREPLANT/ SITE PREPARATION | Glyphosate, MOA 9 Various brands and formulations | See label | Apply 30 days prior to planting for control of emerged weeds. | 12 | Use to kill strips through vineyard prior to planting. Generic formulations may require the addition of a surfactant. See label for details on controlling specific perennial weeds. |
| PREEMERGENCE | | | | | |
| Annual grasses and small seeded broadleaf weeds | Oryzalin, MOA 3 Surflan 4 AS or Oryzalin | 2 to 4 qt | Newly Planted (once soil has settled after transplanting) and Established Vineyards. | 12 | Oryzalin may be tank mixed with paraquat, glyphosate, or Rely for postemergence weed control. In established vineyards tank mix with simazine for expanded residual control of annual weeds. |
| <i>(continued on next page)</i> | Pendimethalin MOA 3 Prowl H ₂ O | 2 to 4 qt | Newly Planted (once soil has settled after transplanting) and established vineyards. | 12 | In newly planted grapes allow soil to settle after transplanting before applying Prowl. Use only during dormancy (prior to bud swell) when applying around newly planted and 1-year-old vines. In bearing vineyards apply any time after harvest, during winter dormancy in spring, and in season before harvest. Use rate cannot exceed 6.3 quarts/acre per year. Prowl H ₂ O has a 21-day PHI. Tank mix with Zeus Prime, simazine or rimsulfuron for expanded residual control of broadleaf weeds. Apply in combination with paraquat, glyphosate, or glufosinate for non-selective POST weed control. |

| Weed/Timing | Material | Amount of Formulation per Acre | Crop Age Restrictions | REI (hrs) | Comments |
|--|--|-------------------------------------|--|-----------|---|
| PREEMERGENCE | | | | | |
| Annual broadleaf weeds | Isoxaben, MOA 21 Trellis SC | 16 to 31 fl. oz | Newly planted and established vineyards | 12 | In newly planted vineyards apply once soil has settled after transplanting. Total use rate per year (from harvest to harvest) cannot exceed 31 fl. oz/A. Trellis SC has a 60 day PHI. For residual control of annual grasses, tank mix with oryzalin or pendimethalin. Tank mix with glyphosate, paraquat, or glufosinate for non-selective POST weed control. |
| Annual weeds and some perennial weeds | Dichlobenil, MOA 20 Casoron 4G Or Casoron 1.4 CS | 100 to 150 lb 1.4 to 2.8 gal | Newly planted (4 wks after transplanting) and established vineyards. | 12 | Apply in January or February for best results. Warm temperatures increase volatilization therefore overhead irrigation may be use for activation when applied in early spring. The Casoron CS formulation should only be used under well-established vines (1 year or older). The Casoron 4G formulation may be used as early as 4 weeks after transplanting young vines. |
| Broadleaf weeds and suppression of yellow nutsedge | Rimsulfuron, MOA 2 Grapple 25 WG Matrix 25 WG Pravin 25 WG Solida 25 WG | 4 oz | Vines established at least 1 year. | 4 | Tank mix with oryzalin, diuron, or simazine to broaden spectrum of residual control. DO NOT apply within 14 days of harvest. Rimsulfuron will provide POST weed control of certain species like horseweed, wild radish, pigweed, chickweed, and henbit. Tank mix with glufosinate, glyphosate, or paraquat for non-selective POST weed control. Tank mixes with glyphosate will provide partial control of yellow nutsedge (2 to 3" tall). |
| Broadleaf weeds and some annual grasses | Diuron, MOA 7 Karmex 80 XP Or Direx 80 DF | 2 to 3 lb | Vines established at least 3 years. | 12 | Heavy rainfall soon after application to grapes planted in soils low in clay and <2% organic matter may result in severe injury and this risk is assumed by the user. Apply with glyphosate, paraquat or glufosinate for postemergence weed control. |

| Weed/Timing | Material | Amount of Formulation per Acre | Crop Age Restrictions | REI (hrs) | Comments |
|---|---|----------------------------------|---|-----------|--|
| PREEMERGENCE | | | | | |
| Broadleaf weeds and some annual grasses | Simazine, MOA 5 Princep 4 L or Princep Cal 90 or various generic formulations | 2 to 4 qt 2.2 to 4.4 lb | Vines established at least 3 years. | 12 | Tank mix with glyphosate, paraquat, or glufosinate for postemergence weed control. The addition of oryzalin (Surflan) or norflurazon (Solicam) or pendimethalin (Prowl H ₂ O) with simazine will extend residual grass control several weeks. |
| Annual broadleaf and grass weeds | Flumioxazin, MOA 14 Chateau 51 SW Tuscany 51 WDG Tuscany SC | 6 to 12 oz 6 to 12 fl. oz | Newly planted and established vineyards | 12 | Apply with hooded or shielded application equipment. Grapes established less than 2 years must be shielded with grow tubes. Flumioxazin may only be used in table grapes after completing harvest and before bud break. Flumioxazin may be applied in vineyards producing grapes used for wine or juice after bud break so long as hooded application equipment is used. DO NOT tank mix with glyphosate after bud break. DO NOT apply more than 6 oz per acre to vines established less than 3 years planted on soils having a sand plus gravel content that exceeds 80%. Flumioxazin formulations have a 60-day PHI. |
| Annual broadleaf and grass weeds (continued) | Indaziflam, MOA 29 Alion 1.67 SC | 5 oz | Vines established at least 5 years | 12 | DO NOT apply to grapes grown in Georgia or Florida. Alion may be used on soils having a texture of sandy loam or finer and less than 20% gravel content. Tank mix with paraquat, glyphosate, or glufosinate for non-selective POST weed control. DO NOT exceed 5 oz of Alion per acre within a 12-month period. If making more than one application per year allow at least 90 days between applications. Tank-mix with glufosinate, glyphosate or paraquat for non-selective POST weed control. |

| Weed/Timing | Material | Amount of Formulation per Acre | Crop Age Restrictions | REI (hrs) | Comments |
|---|---|--------------------------------|--|-----------|--|
| PREEMERGENCE | | | | | |
| Annual broadleaf, some grass weeds, and yellow nutsedge | Carfentrazone + Sulfentrazone, MOA14 Zeus Prime | 7.7 to 15.2 fl. oz | Vines established 2 years or more | 12 | DO NOT allow spray solution to contact green bark or desirable foliage. Zeus Prime XC should be tank mixed with oryzaline for broadspectrum residual control of annual grass weeds. Sequential applications can be made so long as the herbicide strip width is 50% or less of the vineyard floor. Allow 60 days between applications. DO NOT tank mix with flumioxazin. A ½ inch of rainfall is needed within 14 days of application to insure herbicide activation. Tank mix with glyphosate, glufosinate or paraquat for non-selective POST weed control. |
| POSTEMERGENCE DIRECTED | | | | | |
| POSTEMERGENCE DIRECTED Non-selective control | Glyphosate, MOA 9 Various Brands and Formulations 4 SL | See Label | Vines established 1 year or more. | 12 | DO NOT allow spray solution to contact green bark, foliage, or suckers. Tank mix with preemergence herbicides for residual control. Do not apply within 14 days of harvest. Generic formulations may require the addition of a surfactant. Refer to label for application directions for hard to control perennial species. |
| <i>(continued on next page)</i> | Glufosinate, MOA 10 Cheetah Lifeline, Reckon 280 Rely 280 Surmise | 48 to 82 oz | Newly planted (shielded) and established vineyards | 12 | Do not allow herbicide to contact desirable foliage or immature, uncalled bark. Apply in a minimum spray volume of 20 gal./A. Do not apply within 14 days of harvest. |

| Weed/Timing | Material | Amount of Formulation per Acre | Crop Age Restrictions | REI (hrs) | Comments |
|--|--|--------------------------------|---|-----------|---|
| POSTEMERGENCE DIRECTED Non-selective control <i>(continued)</i> | Paraquat, MOA 22 Gramoxone Parazone, or Paraquat Concentrate 3SL | 1.7 to 2.7 pt | Newly planted (shielded) to established vineyards | 12 | Do not allow herbicide to contact desirable foliage or immature, uncalled bark. Young vines must be shielded. Apply in a minimum spray volume of 20 gal./A with non-ionic surfactant at 0.25 % v/v (1qt per 100 gal. of spray solution). Please note additional training requirements for applicators. |
| POSTEMERGENCE DIRECTED Certain broadleaf weeds | Carfentrazone MOA 14 Aim | 1 to 2 fl. oz | Vines established 1 year or longer. | 12 | Do not allow herbicide to contact desirable fruit or foliage. The addition of a non-ionic surfactant at 0.25 % v/v (1 qt per 100 gal of solution) or crop oil concentrate at 1% v/v (1gal per 100 gal. of solution) is necessary for optimum herbicide performance. Ammonium sulfate may be used in addition to a non-ionic surfactant, refer to label for details. Aim may be tank mixed with glyphosate or Rely or various preemergence herbicides. For chemical removal of suckers use the maximum rate and refer to label for details. Aim has a 3-day PHI. |
| POSTEMERGENCE Annual and perennial grasses | Clethodim, MOA 1 Select, and others 2EC SelectMax, | 6 to 8 oz 12 to 16 oz | Newly planted or non-bearing vineyards | 12 | Sequential applications are for perennial grasses (bermudagrass or johnsongrass). The addition of a non-ionic surfactant at 0.25 % v/v (1 qt/100 gal. of spray solution) is required. Do not apply within 1 year of harvest. |
| <i>(continued on next page)</i> | Fluazifop, MOA 1 Fusilade DX | 12 to 24 oz | Newly planted and non-bearing vineyards | 12 | Sequential applications will be necessary for perennial grass (bermudagrass, etc.) control. The addition of a non-ionic surfactant (1 qt/100 gal of spray solution) or crop oil concentrate (1 gal./100 gal. of spray solution) is necessary for optimum results. Do not apply within 1 year of harvest. |

| Weed/Timing | Material | Amount of Formulation per Acre | Crop Age Restrictions | REI (hrs) | Comments |
|--|---|--------------------------------|---|-----------|---|
| POSTEMERGENCE Annual and perennial grasses | Sethoxydim, MOA 1 Poast | 1 to 2.5 pt | Newly planted and established vineyards | 12 | Sequential applications will be necessary for perennial grass (bermudagrass, etc.) control. The addition of a non-ionic surfactant (1 qt/100 gal of spray solution) or crop oil concentrate (1 gal./100 gal. of spray solution) is necessary for optimum results. Do not apply within 50 days of harvest. Total use cannot exceed 5 pt/A per year. |

Suggested Herbicide Programs for Grape Vineyards

| Crop Age | Fall | Winter | Spring | Summer |
|---|--|--|--|--|
| Newly Planted | Glyphosate (Pre-Plant to kill weeds in herbicide strip) | | Flumioxazin (Once soil settles after transplanting) | Flumioxazin + Paraquat (June or July); Fusilade, or Poast, or Clethodim (as needed). |
| Vines Established 1 to 2 years or more | Glyphosate (spot treat for perennial weeds) | Glyphosate (Mid March) | | |
| | Glyphosate (spot treat for perennial weeds) | Flumioxazin + glyphosate, paraquat or Glufosinate (mid to late March) | Flumioxazin* + Paraquat or Glufosinate (early June) | Poast (as needed for POST grass control) *See Flumioxazin restrictions for applications made after bud break. |
| | Glyphosate (spot treat for perennial weeds) | Zeus Prime + Prowl H2O (vines est. 2 yrs) + glyphosate, paraquat, or Glufosinate | Zeus Prime + Prowl H2O + glyphosate, paraquat, or Glufosinate | Glyphosate, Paraquat, Glufosinate, or Poast (as needed) |
| | Glyphosate (spot treat for perennial weeds); Flumioxazin + Glufosinate (after harvest) | | Flumioxazin* + Paraquat, or Glufosinate (late May) | Glufosinate or Paraquat or Poast (as needed) *See Flumioxazin restrictions for applications made after bud break. |
| | Glyphosate (spot treat for perennial weeds) | Flumioxazin + Glyphosate (prior to bud break) | | Glufosinate or Paraquat or Poast (as needed) |
| Vines Established at least 3 years or more | Glyphosate (spot treat for perennial weeds) | Glyphosate (mid-March) | Simazine + Prowl H2O + Glyphosate | Paraquat, Glufosinate, or Poast (as needed) |
| | Glyphosate (spot treat for perennial weeds) | Glyphosate (mid-March) | Karmex + Rimsulfuron (Matrix and others) + Glyphosate or Glufosinate | Paraquat, Glufosinate, or Poast (as needed) |
| Vines Established 5 years or more (continued on next page) | Glyphosate (spot treat for perennial weeds); Simazine + Glufosinate or Paraquat | | Alion + Glyphosate or Glufosinate (Late May to early June) | Paraquat, Glufosinate, or Poast (as needed) |

| | | | | |
|--|--|------------------------|--|---|
| Vines Established 5 years or more <i>(continued)</i> | Glyphosate (spot treat for perennial weeds); Flumioxazin + Glufosinate or Paraquat | | Alion + Glyphosate or Glufosinate (late May to early June) | Paraquat, Glufosinate, or Poast (as needed) |
| | Glyphosate (spot treat for perennial weeds) | Glyphosate (mid-March) | Alion + Glyphosate or Glufosinate (early to mid-May) | Paraquat, Glufosinate, or Poast (as needed) |

Weed Response to Vineyard Herbicides

| Herbicides | Annual Grasses | | | | | Annual Broadleaf Weeds | | | | | | | | | | | | | | | Perennial Weeds | | | | | | |
|------------------------|----------------|----------|------------|---------------|------------------|------------------------|------|-----------|--------------------|-------------------|--------|-----------|---------------|---------------------|-------------|---------|--------------|---------|---------------|-----------|-----------------|--------------|-----------|--------------|------------------|------------------|---|
| | Crabgrass | Foxtails | Goosegrass | Panicum, Fall | Ryegrass, Annual | Chickweed | Dock | Galinsoga | Geranium, Carolina | Groundsel, Common | Henbit | Horseweed | Lambsquarters | Mornigglory, Annual | Nightshades | Pigweed | Radish, Wild | Ragweed | Sida, Prickly | Smartweed | Spotted Spurge | Bermudagrass | Dandelion | Johnsongrass | Nutsedge, Yellow | Virginia Creeper | |
| Preemergence | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alion | E | E | E | G | G | E | | E | E | | E | G | E | E | E | E | G | E | G | G | E | N | G | | P | N | |
| Casoron | G | G | G | G | G | G | G | F | G | G | G | G | G | F | F | G | G | G | | G | G | N | G | | N | N | |
| Flumioxazin | E | E | E | G | G | E | | G | G | | E | G | E | E | E | E | G | G | E | G | E | N | G | | N | N | |
| Diuron | G | G | G | F | G | G | | G | F | | G | G | G | G | G | G | G | G | G | G | N | N | N | | N | N | |
| Kerb | G | G | G | G | G | G | | P | | | G | | F | F | F | P | F | F | | F | | | P | | P | N | |
| Rimsulfuron | F | F | P | P | P | G | | | | G | G | E | G | G | F | E | G | F | | | G | | F | | F | | |
| Oryzalin | E | E | E | G | G | G | N | N | | F | F | | E | F | P | E | P | P | P | P | F | N | P | | N | N | |
| Prowl H ₂ O | E | G | G | G | G | G | | | G | | G | | G | F | F | E | G | | | G | G | | | | | | |
| Simazine | F | G | G | F | G | G | | G | F | F | G | G | E | F | G | G | E | G | F | G | P | N | P | | N | N | |
| Zeus Prime | F | F | F | F | F | G | G | G | G | G | G | F | E | E | E | E | E | F | E | E | E | N | | N | E | N | |
| Postemergence | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Aim | N | N | N | N | N | | | | | | | | P | G | E | G | G | F | | | G | | N | N | N | N | N |
| Clethodim | E | E | E | E | E | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | E | N | | N | N |
| Fusilade | G | G | G | G | G | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | E | N | | N | N | |
| Glyphosate | E | E | E | E | E | E | G | G | G | E | F | E | E | G | E | E | G | E | G | F | G | F | G | | F | G | |
| Paraquat | G | G | G | G | G | G | | G | F | F | F | P | G | G | G | G | F | G | G | G | G | P | P | | P | P | |
| Poast | E | E | E | E | G | N | F | N | N | N | N | N | N | N | N | N | N | N | N | N | N | E | N | | N | N | |
| Glufosinate | F | G | G | G | G | G | N | F | F | F | F | E | G | E | G | G | G | G | F | G | G | F | G | | F | P | |

E = excellent, G = good, F = fair, P = poor, N = no activity

Postemergence Control of Bermudagrass and Johnsongrass

Perennial grasses like bermudagrass and Johnsongrass can be controlled with Poast, Fusilade, and clethodim. Successful use of grass-specific herbicides (Poast, Fusilade DX, clethodim) depends on several factors however the most critical is application timing relative to weed growth stage. Application timing varies with grass species and somewhat with the herbicide choice which is outlined in table below. Additional factors influencing the performance of these herbicides on perennial grasses include spray volume and soil moisture. Graminicides are systemic herbicides, they enter the plant and move through the vascular system to their targeted site of action. Systemic herbicides need to be applied in spray volumes that do not exceed 25 gal. of spray solution per acre. Higher volumes dilute the herbicide and may reduce their effectiveness. Weeds free of stress (drought, etc.) also respond best to systemic herbicides because the herbicide moves into plant and through its vascular system more readily. All of these herbicides require a second application for them to be effective. It is important that the second application be timed appropriately and when the weed has regrown from the initial herbicide application. The time between the first and second application can vary depending upon environmental conditions so this requires monitoring in order to get the second application applied timely.

Appropriate Application Time for Perennial Grass Control

| Herbicide | Bermudagrass | | Johnsongrass | |
|-----------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|
| | 1 st Application | 2 nd Application | 1 st Application | 2 nd Application |
| Poast | 6 inches | 4 inches | 25 inches | 12 inches |
| Clethodim | 3-6 inches | 3-6 inches | 12-24 inches | 6-18 inches |
| Fusilade | 4-8 inches | 4-8 inches | 8-16 inches | 6-12 inches |

Refer to product label for spray additive recommendations.

| Wildlife Damage Prevention | |
|--|--|
| Pest/Problem | Management Options |
| <p>Efforts to control birds and other wildlife that damage fruit crops should be focused on the perimeter of the planting first, especially on the side(s) facing favorable wildlife habitat. This is where the first damage will be observed and, in some cases, it may be sufficient to head off the problem. However, don't discontinue monitoring for wildlife damage throughout the planting.</p> | |
| Birds | <p>Crop losses to birds appear to be increasing in small fruit crops. Not only do birds consume fruit, but the damage they cause can result in increased problems with fruit rots and other pests such as bees and yellow jackets. Several different types of birds can cause problems. Robins, starlings and mockingbirds are among the more common ones, but orioles, cedar waxwings and finches may also feed on small fruit crops.</p> <p>Feeding pressure will be heavier in fields that are close to roosting or nesting sites such as woodlands, hedgerows, grassy fields, power lines and individual trees. Birds may feed, fly to these resting sites, and then return to the crop later in the day. While birds can and do fly fairly long distances to feed, the further they have to fly, the more apt they are to not find the fruit crop or to be distracted by another food source. The presence of a pond, creek or other water source nearby is another factor that may lead to increased feeding pressure. Typically, bird damage tends to be more severe in the earlier parts of the growing season, and damage lessens as the season progresses.</p> <p>There are several control techniques which may be of value in decreasing losses to birds. They include visual, auditory and chemical repellents and exclusion (netting). For any method to be successful, it must be instituted before birds establish a feeding pattern, which generally means that they should be in place and operating at the time that color change occurs in the fruit. With the exception of exclusion, no one method should be relied on for control.</p> <p>Auditory repellents Auditory scare devices such as propane cannons, noise makers or distress calls may offer temporary relief for some types of birds. Regardless of which one or ones is/are used, the following points should be considered to attain the best results:</p> <ul style="list-style-type: none"> - Assess the potential for objections to the noise from your neighbors. - Start before birds establish a feeding pattern. - Operate control devices beginning shortly before sunrise and continuing until just after sunset, as early and late in the day may be the most intense feeding times. - Vary the frequency, the direction and the timing in which auditory devices are operated. Propane cannons should not be fired at intervals |

| | |
|------------------------------|--|
| <p>Birds (Continued)</p> | <p>closer than 3 minutes.</p> <ul style="list-style-type: none"> - Consider using more than one type of auditory device and possibly combine them with visual repellents. - If using distress calls, it is essential to identify the type(s) of birds you want to discourage and get distress calls specific to them. - Reinforce the sense of danger by shooting (if allowed). <p>Visual repellents Visual repellents include such things as scare eyes suspended above the crop, mylar tape on the canopy of the crop, aluminum pie pans, plastic owls and plastic snakes. These range from ineffective to moderately effective for a short period of time. Birds will get used to them quickly if they are not moved around or if another type of repellent is not used along with it. Yellow scare eyes suspended above the crop and allowed to move freely have been reported to have some impact on blackbirds, however, robins do not seem to be affected.</p> <p>Chemical repellents Methyl anthranilate is registered as a bird repellent. While it is sometimes advertised as a taste repellent, this is not exactly correct. When sprayed on a crop, it causes an unpleasant sensation in the bird's mouth. Methyl anthranilate is a naturally occurring material used in the food service industry. Early reports have been inconsistent in regards to its effectiveness. It has also been reported to impart an undesirable foxy flavor to certain grape varieties. Methyl anthranilate has a short residual, so frequent reapplication will be necessary to achieve lasting results. Results may vary depending on the type of birds. Combining with another type of deterrent may result in greater effect than when used alone. As with other types of deterrents, applications need to start before birds establish feeding patterns.</p> <p>Exclusion Exclusion (netting) is the only consistently effective method of reducing bird damage. Netting is more expensive than other types of deterrents and can require fair amounts of labor, so it may not be an economically viable alternative in all situations. Nets are either laid on the canopy of the crop or suspended from a framework over the crop. The fruiting area of the plant needs to be completely protected. Birds will enter the canopy of the plant from below the net if it is open under the plant. If used with care, nets can be maintained for use over several years. For crops requiring multiple harvests such as blueberry, suspending the netting over the crop and around the sides of the field will allow easier access to the crop. If nets are placed directly on the crop canopy, birds can perch on it and feed on berries below them.</p> <p>Wild turkeys are becoming more of a problem in many areas of the country. While there is no doubt that they do consume some fruit, some research has shown that the turkeys are often after insects instead of the fruit. They do not appear to like loud and/or distressing sounds. While netting will work, turkeys can tear holes in it for access to the fruit.</p> |
| <p>Deer</p> | <p>Deer can damage small fruit plantings by foraging on succulent new growth during the growing season or by eating fruit. In fall, bucks can damage plants by rubbing. This is more of a problem in tree fruits than small fruits. Deer can also puncture plastic mulch and possibly the irrigation tape underneath, resulting in loss of weed control. Deer numbers are increasing and incidents of deer damaging crops are also increasing. Deer populations vary from year to year as a result of weather conditions, food supply and possibly hunting pressure.</p> <p>Locating the planting away from favorable habitat for deer will help to lessen losses. However, this is not always possible. Several control options do exist. Determining which one or ones to use depends on the deer population, availability of other food sources, location of favorable habitat, the duration for which protection is needed, and the value of the crop to be protected.</p> |

| | |
|-----------------------------|--|
| <p>Deer (Continued)</p> | <p>Repellents Both taste and smell repellents exist. Smell repellents include commercially available products or materials such as tankage, blood, putrified egg solids, certain soaps and human hair. Repellants will not provide long-term control and will not provide control when populations are high or alternate food sources are scarce.</p> <p>Exclusion Exclusion (fencing) is the only truly effective long-term control for deer damage prevention. Fences can be electrified or not. Deer will try to go under a fence, through a fence, or over it. For non-electrified fences, the lowest wire needs to be within 10 inches or less of the lowest point in the ground around the fruit crop planting and tight enough to prevent deer from pushing under it. Do not neglect ditches or other low spots in the ground around the field, because the deer will find them. The fence needs to be at least 8 feet high or higher, as deer can easily clear this height. Wire mesh fences are more desirable than multiple strands of barbed wire.</p> <p>For electric fences, several different designs have been used and, under certain conditions, each can be effective. The simplest and least expensive electric fence uses a single high-tensile wire at about 30 inches above ground level. A solar charger can be used if access to electricity is not an option. Peanut butter can either be smeared on the wire or on aluminum foil strips which are then draped over the wire. Plastic flagging may also be tied to the fence to make it more visible to the deer. Deer are curious animals and will investigate the fence if they are not being chased. Touching the fence results in getting shocked and turning the deer away from the field being protected. The single-wire, baited fence is relatively inexpensive, easy to construct and often adequate to protect the crop. With high deer populations, when available alternate food sources are scarce or when deer have already established a feeding pattern in the area being protected, this fence may not be adequate.</p> <p>More substantial electric fences for deer control have multiple wires with the alternate wires being electrified. One design uses 5 wires and is constructed at a 45-degree angle facing away from the area to be protected. The bottom wire is within 10 inches of the ground and is electrified to keep deer from going under the fence. The middle wire is also electrified to prevent deer from going through the fence and the top wire, which may be only about 5 feet above ground is electrified to keep deer from going over the fence. A fence constructed in this manner presents a barrier to the deer that has height and depth, a combination that generally will discourage the deer from trying to enter the field. Poly Tape electric fence often used to contain cattle and horses works well for deer fences.</p> <p>Numerous other fence designs exist including a non-electrified mesh fence with a hot wire on top. If electric fences are used, it is important to keep weeds, grasses and other materials away from the fence to prevent it from shorting out and to increase its visibility.</p> |
|-----------------------------|--|

The University of Georgia and Ft. Valley State University, the U.S. Department of Agriculture and counties of the state cooperating. The Cooperative Extension Service offers educational programs, assistance and materials to all people without regard to race, color, national origin, age, sex or disability.

An Equal Opportunity Employer/Affirmative Action Organization Committed to a Diverse Work Force

extension.uga.edu

Annual Publication 126-1

March 2022

Published by the University of Georgia in cooperation with Fort Valley State University, the U.S. Department of Agriculture, and counties of the state. For more information, contact your local UGA Cooperative Extension office.
The University of Georgia College of Agricultural and Environmental Sciences (working cooperatively with Fort Valley State University, the U.S. Department of Agriculture, and the counties of Georgia) offers its educational programs, assistance, and materials to all people without regard to race, color, religion, sex, national origin, disability, gender identity, sexual orientation or protected veteran status and is an Equal Opportunity, Affirmative Action organization.