Michel Kohl, Wildlife Specialist

NOTE: For recommendations on the use of chemicals to control vertebrate pests, contact your Cooperative Extension Service agent. It is necessary to obtain a permit prior to killing protected animals in Georgia. Get a permit by writing the Wildlife Resources Office of the Georgia Department of Natural Resources, 2070 U.S. Highway 278, S.E., Social Circle, Georgia 30279; or call 706-557-3305.

In the following section, non-game animals include pocket gophers, chipmunks, bats, amphibians, and reptiles. A permit is also required to take furbearing

animals or game animals out of season. A current hunting or trapping license is required to take game or furbearing animals in season.

To begin the permit process for taking protected birds, call USDA Wildlife Services at 706-546-5637. For permits to kill other protected animals, call the Wildlife Resources office of the Georgia Department of Natural Resources 706-557-3304.

MAMMALS

SPECIES	HABITAT MODIFICATION	EXCLUSION	FRIGHTENING	REPELLENTS	TOXICANTS/ FUMIGANTS	TRAPPING	COMMENTS
Armadillos	No habitat modification options available; armadillo are habitat generalists.	Fencing may be effective for gardens and high-value land areas (flower beds, etc.).	Not effective	None	None are registered	Possibly effective if done properly. See armadillo publication: https://resources. ipmcenters.org/resource. cfm?rid=17253	Lethal control if legal and safe. Armadillo are not protected in Georgia.
Beaver	Eliminate food, trees, woody vegetation where feasible. Continually destroy dams. Install water control device (Clemson beaver pond leveler).	Fence small areas along stream and yard. Fence around valuable trees.	Continual destruction of lodges and dams	None are registered	None are registered	Very effective if done by a trained Wildlife Control Expert.	None Beaver are not protected in Georgia.
Chipmunk	Store food such as bird seed and dog food in rodent-proof container. Ground cover, shrubs, and wood piles should not be located adjacent to struc- tural foundation.	Seal all cracks and holes in foundations. ¹ /4-inch mesh hardware cloth buried 1–2 feet deep to exclude from gardens and flower beds.	Not effective	Taste repellents containing Bitrex or Thiram applied to seeds, bulbs, or vegetation not meant for human consumption.	None are registered	Very effective; use standard mouse or rat snap traps baited with peanut butter.	None
Coyote	Destroy dens if possible. Reduce habitat for mice and other food items.	Fence pastures and yards to reduce danger to livestock and pets. Keep pets inside, especially at night.	Guard dogs can be effective for livestock.	None shown to be consis- tently effective	Available to livestock pro- ducers and Wildlife Control Experts; generally not available to homeowners/ gardeners.	Can be very effective but requires considerable knowledge; best left to Wildlife Control Experts. Georgia DNR maintains a listed of licensed Nuisance Wildlife Trappers.	Coyote are not protected in Georgia.

SPECIES	HABITAT MODIFICATION	EXCLUSION	FRIGHTENING	REPELLENTS	TOXICANTS/ FUMIGANTS	TRAPPING	COMMENTS
Deer	Difficult; habitat modi- fication is not generally recommended. Plant deer-resistant plants Harvest crops early. Lure crops may draw deer away from valuable crops.	Fences—up to 8 feet tall; less if with dog. Electric fences can be effective. Individual tree protector; fence or pipe to prevent antler rubs. Many effective fence designs for high value crops such as fruit, berry, & grapes.	May provide some limited protection in certain situations.	Many are available; some commercial; some "home" remedies. Effectiveness varies with ingredients and timing; start applying prior to damage; high density deer population will not likely be repelled. Taste and odor repellents are available; beware of advertising. Sound repellents are generally ineffective.	None	Not available to landowners	Contraception is not legal in Georgia. Hunting is best in most sit- uations but not allowable in some instances. Commercial control at night with licensed professionals.
Gophers or Pocket Gophers	No habitat modification options available.	Limited efficacy for small gardens. Not practical for fields or landscapes.	Not effective	None	Some are registered. Check Georgia Department of Agriculture pesticide data- base for current list.	Difficult to trap– requires expertise and special traps.	Not effective except with registered toxicant.
Gray Squirrel or Flying Squirrel	Trim trees away from house. Remove diseased trees or trees with cavities. Remove bird feeders near house or only use squirel-resistent bird feeders.	Sheet metal bands on isolated trees to prevent climbing in order to pro- tect developing nut crop. Seal all openings to build- ings, especially around chimney, eaves, and soffits. Install excluder in attics to allow squirrel to escape and prevent re-entry.	Not effective	Some products containing hot sauce (capsaicin) and other distasteful com- pounds may work (may also be used with deer).	None are registered	Can be very effective but should only be done by qualified Wildlife Control Expert. Permits may be required.	On bird feeders suspended by rope, place rope through plastic conduit to reduce climbing. For squirrel feeders hung on poles, use squirrel baf- fles to prevent climbing.
Hogs— Feral Hogs, Feral Swine, Wild Hogs, Wild Pigs, Wild Swine	Difficult	Fencing may be effective for small areas.	May provide limited pro- tection in some instances.	None. There is no such thing as pig repellent. Mothballs are not labeled for use.	None—NO legal toxicant is available.	Trapping is the best method of control. Effective trapping requires training or knowledge of methods. Very effective when prop- erly applied. Use ONLY large corral style traps.	Shooting or hunting offers limited control. Check all game regulations. Hunting, especially dog hunting, may make the problem worse. Relocating or transporting live wild pigs is illegal and punishable by strict penalties.

MAMMALS & BIRDS

SPECIES	HABITAT MODIFICATION	EXCLUSION	FRIGHTENING	REPELLENTS	TOXICANTS/ FUMIGANTS	TRAPPING	COMMENTS
Moles	Reduce soil moisture and food sources by reducing watering. Cautious use of lawn pesticides to treat for insects and grubs	May be cost effective in very small area like flower bed or garden. ¼-inch hardware cloth or sheet metal.	Not effective	None are registered	Several products (poisons) are available for placing in the tunnel or hole.	Difficult but can be effective in some situations.	None
Voles	Eliminate ground cover such as grass and weeds. Remove mulch from base of trees and plants. Soil cultivation to destroy burrows, tunnels, and ground cover.	Recommended to protect trees, ornamentals, and gardens. Use ¼-inch mesh hardware cloth buried 1–2 feet deep.	Not effective	None proven effective	Anticoagulants, such as rat poisons. Always use caution around pets and children.	Very effective using standard rat or mouse traps baited with peanut butter.	None

BIRDS

SPECIES	PROBLEM STATEMENT	TOXICANTS/ FUMIGANTS	PRECAUTIONS	COMMENTS
Birds, general	Nesting in rafters of pole barns; pecking holes in siding; roosting; eating fruit.	No toxicants or fumigants registered.		Recommendations are for physical exclusion with fence, net, or barrier. Brand name: Bird-X* netting, Nix-A-Lite*, sticky caulking. Grapes and berries—fencing with bird netting; scare devices. Lasers may also deter birds, but may be expensive.
Canada Goose	Droppings on lawn, playground, drive- way; eating grass; nesting.	Methyl anthranilate	Follow label directions.	Goose repellent made from Concord grapes. Expensive. Must reapply after mowing, rain or irrigation. Addling (shaking) eggs or coating with oil may be allowed—contact Wildlife Specialist or USDA APHIS Wildlife Services. Destroy nests with permints; harass with dogs; trap and remove during flightless period.
Woodpeckers	Pecking holes in siding and wooden decks—often in search of food or nesting sites.	No toxicants or fumigants		Physical exclusion with netting; harassment with shiny flagging often called "Mylar Tape". All birds are federally protected—cannot be killed. Permits for persistent problems are available—Contact Wildlife Specialist or USDA APHIS Wildlife Services.

REPELLENTS FOR DEER, RABBITS, AND OTHER MAMMALS

These materials may give temporary protection. Repellents are most likely to produce satisfactory results if protection is needed for a week or two. If longer protection is required, fencing is often a more effective treatment. Re-treatment may be necessary, especially after rain. Repellents are most effective when applied before damage starts. Local density of animals may negate efficacy of any repellent treatment. Some repellents are contact- and water-soluble, others are systemic and therefore require less frequent application. Read and follow all label instructions. Most commercial repellents are **not labeled** for use on food crops. Contact wildlife specialist for additional products.

PROBLEM OR LOCATION	CHEMICAL & FORMULATION	METHOD AND RATE	TARGET SPECIES	COMMENTS
Browsing on conifer seedlings, ornamentals, hardwoods	Putrescent Whole Egg solids. Can be homemade.		Deer	A taste repellent. Not for use on edible plants. Brands or product names: Deer away, BGR.
	Liquid 15% Powder 36%	Apply to vegetation—spray, dip, or shake as per label.	Deer	
Fruit trees, field crops, ornamen- tals, nursery stock, vegetables, non-crop areas	Ammonium soaps of higher fatty acids	Dilute as per label. Spray on crops to be protected.	Deer, Rabbit	Re-treat after rain or as necessary.
	Liquid 15%	May work as border treatment. Label may include use as paint for stems and trunks.	Deer	Brand or product names: Hinder Rabbit
	Other taste repellents: Mint Oil, Garlic Oil, Capsaicin, Bitrex®		Deer, Rabbit	Re-treat after rain or as necessary. Apply before damage is noticed. Follow label instructions–some products are not labeled for edible crops. Brand or product names: Tree-Guard, Liquid Fence, Deer-Off, Phantskydd, This-1-Works*
	Order Repellents: Milorganite®	Apply as per label	Deer, Rabbit	Re-treat as necessary. Most home odor repellents are not-effective. Milorganite [®] is a mild organic fertilizer that is very effective if used properly.
Gnawing on wood siding, decks, fence posts	benzyidlethyl-methyl-ammonium- saccharide and thymol	Apply as per label. Do not apply indoors. Avoid contact with water or food sources. Do not mix with other chemicals.	Rats, Squirrels, Skunks, Rabbits	Manufacturer claims of effectiveness may need further testing. May repel woodpeckers. Brand name: Ro-Pel*

REPTILES AND AMPHIBIANS

There are no intoxicants or fumigants registered for reptiles and amphibians. Commercial snake repellents are available but none have been shown to be consistently effective. Snakes lack a sense of smell and do not hear sound, so repellents are of questionable efficacy. Nearly all reptiles except venomous (poisonous) snakes are protected. Amphibians enjoy less legal protection. Insects like ticks and chiggers can be repelled with products containing DEET. There is no such thing as snake repellent. Mothballs (napthalene) are not labeled as a vertebrate repellent. Use of mothballs for snakes (or anything else) is illegal and ineffective.

ANIMAL	PROBLEM OR LOCATION	CHEMICAL AND FORMULATION	METHOD AND RATE	COMMENTS
Reptiles: snakes, lizards, turtles, and alligators	Yard, in home, barns, and outbuildings	No toxicants or fumigants registered.	Not Applicable	No venomous lizards in Georgia. Six of 40 species of snakes are venomous. All harmless (non-venomous) snakes protected. Call DNR. Do not kill. If dead, specimen should be frozen. Do not damage head. Alligators are protected in Georgia. Contact Georgia Wildlife Resources biologists for alligator problems. Use traps to remove nuisance turtles. Do not use mothballs or "Snake-Away" type products as they do not work.
Amphibians: frogs, toads, salamanders, and their tadpoles	Animals getting into swim- ming pools, houses	None		No chemical controls currently available. Hand removal with a dip net is recommended. Keeping sticks or rocks in small garden pools will allow animal to climb out.