

SMALL GRAIN INSECT CONTROL

David Buntin, Research/Extension Entomologist

GRAIN PRODUCTION OF WHEAT, TRITICALE, BARLEY, OATS, AND RYE

CROP/PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days) H-harvest grain G-grazing, hay	TREATMENT THRESHOLDS AND REMARKS
Insecticides and products listed are for use on all cereal grain crops for grain production including wheat, triticale, barley, oats, and rye, except where listed or noted in the insecticide column and remarks section. Products labeled for use only on wheat or wheat and triticale are <i>dimethoate</i> , Fastac CS, and Tombstone. ► ALL PRODUCTS LISTED IN THIS SECTION HAVE A BEE HAZARD RESTRICTION, EXCEPT FOR <i>CHLORANTRANILIPROLE</i> (CORAGEN, PREVATHON).						
Aphids	SEED TREATMENTS					Inspect fields 25–35 days after planting, full tiller, and heading. Yield-reducing transmission of Barley Yellow Dwarf virus can occur during first two periods; transmission at heading is too late to reduce yield. Aphid treatment thresholds are: • Seedlings (2/row ft) • 6–10 inch plants (6/row ft) • Stem elongation (2/stem) • Flag leaf (5/flag) • Heading (10/head to include flag) • Soft/Hard Dough stages (Do not treat) NOTE: OP insecticides, such as <i>dimethoate</i> , control aphids but are not effective at suppressing Barley Yellow Dwarf disease. NOTE: At labeled rates, Gaucho XT and Enhance AW also contain two fungicides. NOTE: Cruiser 5FS is available as a commercial seed treatment. Rates of CruiserMaxx Cereals and CruiserMaxx Vibrance alone are too low for effective aphid control. NOTE: NipsIt Inside may be sold as NipsIt Suite Cereals CVR that includes 2 fungicides. NOTE: Warrior II Zeon will replace Karate Zeon. NOTE: Transform wheat, triticale and barley only.
	<i>imidacloprid</i> Gaucho 600, Attendant 600 Access Gaucho XT Enhance AW Warden Cereals HR	4A	0.8 fl oz/100 lb seed 3.4 fl oz/100 lb seed 4 fl oz/100 lb 5–8 fl oz/100 lb	0.03 lb/100 lb seed 0.03 lb/100 lb seed 0.05 lb/100 lb seed 0.05–0.08 lb/100 lb seed	12 H/ 45 D	
	<i>thiamethoxam</i> Cruiser 5FS Cruiser Maxx Cereals Plus Cruiser 5FS	4A	1 fl oz/100 lb seed 5 fl oz/100 lb seed plus 0.5 fl oz/100 lb seed	0.04 lb/100 lb seed 0.04 lb/100 lb seed (total)	12 H/ 45 D	
	<i>clothianidin</i> NipsIt Inside	4A	0.75–1.79 fl oz/100 lb seed		12 H/ Not listed	
	FOLIAR TREATMENTS					
	<i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	2.4 fl oz	0.019	12 H/ H-30 D G-3 D	
	<i>dimethoate</i> (wheat only) Dimethoate 4EC, 400 Dimethoate 2.67EC	1B	0.5–0.75 pt 0.75–1 pt	0.25–0.375	48 H/ H-35 D G-35 D	
	<i>flupyradifurone</i> Sivanto Prime	4D	7–10 fl oz	0.09125–0.13	4 H/ H-21 D G-7 D	
	<i>gamma cyhalothrin</i> Declare 1.25 Proaxis 0.5	3A	1.54 fl oz 3.84 fl oz	0.015	24 H/ H-30 D G-7 D	
	<i>lambda cyhalothrin</i> Warrior II Zeon 2.08, Silencer, Lambda, others 1	3A	1.28–1.92 fl oz 2.56–3.84 fl oz	0.02–0.03	24 H/ H-30 D G-7 D	
<i>lambda cyhalothrin</i> + <i>chlorantraniliprole</i> Besiege	3 + 28	5–10 fl oz/A	0.02–0.03 + 0.04–0.06	24 H/ H-30 D G-7 D		
<i>sulfoxaflor</i> Transform 50WG	4C	0.75–1.5 oz	0.023–0.046	24 H/ H-14 D G-7 D		

SMALL GRAIN INSECT CONTROL

SMALL GRAIN

CROP/PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days) H-harvest grain G-grazing, hay	TREATMENT THRESHOLDS AND REMARKS
Armyworm True armyworm Fall armyworm, Beet armyworm, Yellowstriped armyworm, and Cutworms	<i>alpha-cypermethrin</i> Fastac CS 0.83 (wheat, triticale only)	3A	1.8–3.8 fl oz	0.012–0.025	14 H/ H-14 D G-14 D	True armyworm usually infests wheat in late winter and spring at the boot/head stage. Treat when larval numbers exceed 4 larvae/sq ft before pollen shed and 8 larvae/sq ft after pollen shed.
	<i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1.8–2.4 fl oz	0.014–0.019	12 H/ H-30 D G-3 D	
	<i>cyfluthrin</i> (wheat only) Tombstone Tombstone Helios 2	3A	1.8–2.4 fl oz	0.028–0.038	12 H/ H-30 D G-3 D	Fall armyworm, beet armyworm, yellowstriped armyworm and cutworm infestations usually occur in the fall on seedling plants. Treat when larval populations of any one or any combination of these insects exceed 3 larvae (½ inch long or larger)/sq ft.
	<i>chlorantraniliprole</i> Coragen 1.67SC Prevathon 0.43 Vantacor	28	3.5–5 fl oz 14–20 fl oz 1.2–2.5 fl oz	0.045–0.065 0.047–0.067 0.047–0.098	4 H/ H-1 D G-1 D	
	<i>gamma cyhalothrin</i> Declare 1.25 Proaxis 0.5	3A	1.28–1.54 fl oz 3.2–3.84 fl oz	0.0125–0.015	24 H/ 30 D G-7 D	NOTE: Baythroid and Tombstone products are effective against small larvae only.
	<i>lambda cyhalothrin</i> Warrior II Zeon 2.08 Silencer, Lambda, others	3A	1.6–1.92 fl oz 3.2–3.84 fl oz	0.025–0.03	24 H/ H-30 D G-7 D	NOTE: Fastac CS and Tombstone products are not labeled for use on barley, oats, or rye.
	<i>lambda cyhalothrin + chlorantraniliprole</i> Besiege	3 + 28	5–10 fl oz	0.02–0.03 + 0.04–0.06	24 H/ H-30 D G-7 D	
	<i>spinosad</i> Blackhawk (36%)	5	1.7–3.5 oz	0.038–0.075	4 H/ H-21 D G-3 D	
	<i>spinetoram</i> Radiant 1SC	5	3–6 fl oz	0.0234–0.0469	4 H/ H-21 D G-3 D	
<i>zeta-cypermethrin</i> Mustang Maxx, Respect 0.8EC	3A	3.2–4 fl oz	0.02–0.025	12 H/ H-14 D G-3 D		
Cereal Leaf Beetle	<i>alpha-cypermethrin</i> Fastac CS 0.83 (wheat, triticale only)	3A	1.8–3.8 fl oz	0.012–0.025	12 H/ H-14 D G-14 D	Treat when an average of 1 larva and adult/4 stems are found. Warrior II (formerly Karate), Declare and similar products can be applied at 50% egg hatch. Other materials should not be applied until after 90% egg hatch.
	<i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1–1.8 fl oz	0.008–0.014	12 H/ H-30 D G-3 D	
	<i>cyfluthrin</i> (wheat only) Tombstone Tombstone Helios 2	3A	1–1.8 fl oz	0.016–0.028	12 H/ H-30 D G-3 D	NOTE: <i>lambda/gamma cyhalothrin</i> products applied for cereal leaf beetle also provide aphid control for the remainder of the season
	<i>gamma cyhalothrin</i> Declare 1.25 Proaxis 0.5	3A	0.77–1.54 fl oz 2.56–3.84 fl oz	0.0075–0.015 0.01–0.015	24 H/ H-30 D G-7 D	NOTE: Fastac CS, Respect, and Tombstone products are not labeled for use on barley, oats, and rye.

CROP/PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days) H-harvest grain G-grazing, hay	TREATMENT THRESHOLDS AND REMARKS
Cereal Leaf Beetle (continued)	<i>lambda cyhalothrin</i> Warrior II Zeon 2.08 Silencer, Lambda, others	3A	1.28–1.92 fl oz 2.56–3.84 fl oz	0.02–0.03	24 H/ H-30 D G-7 D	Treat when an average of 1 larva and adult/4 stems are found. Warrior II (formerly Karate), Declare and similar products can be applied at 50% egg hatch. Other materials should not be applied until after 90% egg hatch. NOTE: lambda/gamma cyhalothrin products applied for cereal leaf beetle also provide aphid control for the remainder of the season NOTE: Fastac CS, Respect, and Tombstone products are not labeled for use on barley, oats, and rye.
	<i>lambda cyhalothrin + chlorantraniliprole</i> Besiege	3 + 28	5–10 fl oz	0.02–0.03 + 0.04–0.06	24 H/ H-30 D G-7 D	
	<i>malathion</i> Malathion 57EC, 5EC Malathion 8EC	1B	1.5 pt 1 pt	0.94 1	12 H/ H-7 D G-7 D	
	<i>zeta-cypermethrin</i> Mustang Maxx, Respect 0.8EC	3A	2.6–3.2 fl oz	0.015–0.02	12 H/ H-14 D G-14 D	
Grasshoppers	<i>alpha-cypermethrin</i> Fastac CS 0.83 (wheat, triticale only)	3A	3.2–3.8 fl oz	0.020–0.025	12 H/ H-14 D G-14 D	Treat when grasshoppers are causing excessive (greater than 50%) defoliation. NOTE: Fastac CS, Respect, and Tombstone are not labeled for use on barley, oats, and rye. NOTE: Prevathon and Vantacor, for best results add, methylated seed oil (MSO) at 1 gallon per 100 gallons of spray volume (1% v/v).
	<i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1.8–2.4 fl oz	0.014–0.019	12 H/ H-30 D G-3 D	
	<i>cyfluthrin</i> (wheat only) Tombstone Tombstone Helios 2	3A	1.8–2.4 fl oz	0.028–0.038	12 H/ H-30 D G-3 D	
	<i>chlorantraniliprole</i> Coragen 1.67SC Prevathon 0.43 Vantacor	28	3.5–5 fl oz 14–20 fl oz 0.7–1.7 fl oz	0.045–0.065 0.047–0.067 0.027–0.066	4 H/ H-1 D G-1 D	
	<i>gamma cyhalothrin</i> Declare 1.25 Proaxis 0.5	3A	1.02–1.54 fl oz 2.56–3.84 fl oz	0.01–0.015	24 H/ H-30 D G-7 D	
	<i>lambda cyhalothrin</i> Warrior II Zeon 2.08 Silencer, Lambda, others	3A	1.28–1.92 fl oz 2.56–3.84 fl oz	0.02–0.03	24 H/ H-30 D G-3 D	
	<i>lambda cyhalothrin + chlorantraniliprole</i> Besiege	3 + 28	5–10 fl oz	0.02–0.03 + 0.04–0.06	24 H/ H-30 D G-7 D	
	<i>malathion</i> Malathion 57EC, 5EC Malathion 8EC	1B	1.5 pt 1 pt	0.94 1	12 H/ H-7 D G-7 D	
	<i>zeta-cypermethrin</i> Mustang Maxx, Respect 0.8EC	3A	3.2–4 fl oz	0.02–0.025	12 H/ H-14 D G-14 D	

SMALL GRAIN INSECT CONTROL

CROP/PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days) H-harvest grain G-grazing, hay	TREATMENT THRESHOLDS AND REMARKS
Chinch bug	<i>alpha-cypermethrin</i> Fastac CS 0.83 (wheat, triticale only)	3A	3.8 fl oz	0.025	12 H/ H-14 D G-14 D	<p>Treat during the seedling stage when an average of 1 adult/2 plants are found. On larger plants treat when 1 adult/stem or 50% of the plants are infested.</p> <p>Gaucho, Attendant, and Cruiser seed treatments may provide control for a few weeks after planting. Chinch bugs are difficult to control in headed wheat.</p> <p>NOTE: Fastac CS, Respect, and Tombstone are not labeled for use on barley, oats, and rye.</p>
	<i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	2.4 fl oz	0.019	12 H/ H-30 D G-3 D	
	<i>cyfluthrin</i> (wheat only) Tombstone Tombstone Helios 2	3A	2.4 fl oz	0.038	12 H/ H-30 D G-3 D	
	<i>gamma cyhalothrin</i> Declare 1.25 Proaxis 0.5	3A	1.54 fl oz 3.84 fl oz	0.015	24 H/ H-30 D G-7 D	
	<i>lambda cyhalothrin</i> Warrior II Zeon 2.08, Silencer, Lambda, others	3A	1.92 fl oz 3.84 fl oz	0.03	24 H/ H-30 D G-7 D	
	<i>lambda cyhalothrin</i> + <i>chlorantraniliprole</i> Besiege	3 + 28	5–10 fl oz	0.02–0.03 + 0.04–0.06	24 H/ H-30 D G-7 D	
	<i>zeta-cypermethrin</i> Mustang Maxx, Respect 0.8EC	3A	4 fl oz	0.025	12 H/ H-14 D G-14 D	
Hessian fly	SEED TREATMENTS					<p>Plant Hessian fly-resistant wheat varieties.</p> <p>Systemic insecticide seed treatments are recommended for susceptible varieties. Systemic seed treatments may need highest rates for effective suppression. Gaucho XT alone may not provide effective control. Rate of CruiserMaxx Cereals or Cruiser Maxx Vibrance alone is too low for effective Hessian fly control.</p> <p>NOTE: Barley is tolerant. Damage only occurs under severe infestations. Rye is fairly resistant but can be damaged with large infestations.</p> <p>NOTE: Apply Warrior II (formerly Karate) or Declare when adults are actively laying eggs. Apply based on egg sampling for best results.</p>
	<i>imidacloprid</i> Gaucho 600, Attendant 600 Acess Gaucho XT plus Gaucho 600	4A	1.6–2.4 fl oz/100 lb seed 3.4 fl oz/100 lb seed plus 1 fl oz/100 lb seed	0.0625–0.094 lb/ 100 lb seed Combined: 0.0675lb/ 100 lb seed	12 H/ 45 D	
	<i>thiamethoxam</i> Cruiser 5FS Cruiser Maxx Cereals Plus Cruiser 5FS	4A	1.33 fl oz/100 lb seed 5 fl oz/100 lb seed plus 0.5 fl oz/100 lb seed	0.06 lb/100 lb seed 0.06 lb/100 lb seed (total)	12 H/ 45 D	
	<i>clothianidin</i> NipsIt Inside	4A	1.79 fl oz/ 100 lb seed	—	12 H/ Not listed	
	FOLIAR TREATMENTS					
	<i>gamma cyhalothrin</i> Declare 1.25 Proaxis 0.5	3A	1.54 fl oz 3.84 fl oz	0.015	24 H/ H-30 D G-7 D	
	<i>lambda cyhalothrin</i> Warrior II Zeon 2.08, Silencer, Lambda, others	3A	1.92 fl oz 3.84 fl oz	0.03	24 H/ H-30 D G-7 D	

CROP/PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days) H-harvest grain G-grazing, hay	TREATMENT THRESHOLDS AND REMARKS
Hessian fly (continued)	FOLIAR TREATMENTS (continued)					
	<i>lambda cyhalothrin</i> + <i>chlorantraniliprole</i> Besiege	3 + 28	5–10 fl oz	0.02–0.03 + 0.04–0.06	24 H/ H-30 D G-7 D	
Mites/Winter grain mite	<i>gamma cyhalothrin</i> Declare 1.25 Proaxis 0.5	3A	1.54 fl oz 3.84 fl oz	0.015	24 H/ H-30 D G-2 D	Treat when infestations are causing leaf discoloration in large areas of a field and mites are present. Infestations are usually associated with application of cattle manure or chicken litter. NOTE: Warrior II (formerly Karate) and Declare only provide mite suppression.
	<i>lambda cyhalothrin</i> Warrior II Zeon 2.08 Silencer, Lambda, others	3A	1.92 fl oz 3.84 fl oz	0.03	24 H/ H-30 D G-7 D	
	<i>lambda cyhalothrin</i> + <i>chlorantraniliprole</i> Besiege	3 + 28	5–10 fl oz	0.02–0.03 + 0.04–0.06	24 H/ H-30 D G-7 D	
Stink bugs	<i>alpha-cypermethrin</i> Fastac CS 0.83 (wheat, triticale only)	3A	3.2–3.8 fl oz	0.020–0.025	12 H/ H-14 D G-14 D	Stink bugs rarely reach levels needing control in wheat. Treat if stinks bugs exceed 1 bug/sq ft at milk stage. Control during medium-hard dough is rarely justified. NOTE: Fastac CS, Respect, and Tombstone are not labeled for use on barley, oats, and rye.
	<i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1.8–2.4 fl oz	0.014–0.019	12 H/ H-30 D G-7 D	
	<i>cyfluthrin</i> (wheat only) Tombstone Tombstone Helios 2	3A	1.8–2.4 fl oz	0.028–0.038	12 H/ H-30 D G-3 D	
	<i>gamma cyhalothrin</i> Declare 1.25 Proaxis 0.5	3A	1.02–1.54 fl oz 2.56–3.84 fl oz	0.01–0.015	24 H/ H-30 D G-7 D	
	<i>lambda cyhalothrin</i> Warrior II Zeon 2.08 Silencer, Lambda, others	3A	1.28–1.92 fl oz 2.56–3.84 fl oz	0.02–0.03	24 H/ H-30 D G-7 D	
	<i>zeta-cypermethrin</i> Mustang Maxx, Respect 0.8EC	3A	3.2–4 fl oz	0.02–0.025	12 H/ H-14 D G-14 D	

SMALL GRAIN FUNGICIDE SEED TREATMENTS

Alfredo Martinez-Espinoza, Extension Plant Pathologist

WHEAT, OATS, BARLEY, RYE, SORGHUM

FUNGICIDE	CROP	RATE/100 LB SEED	REMARKS
<i>azoxystrobin</i> Dynasty	Wheat and Barley	0.153–0.882 fl oz	For protection against common bunt and partial control of dwarf bunt. Where appropriate use in combination with Dividend extreme.
<i>captan</i> Captan 400	Wheat, Barley, Oats, Rye	See label	Controls seedling blights. Does not control smuts.
<i>carboxin</i> + <i>captan</i> Enhance	Wheat, Barley, Oats	4 oz	Controls loose smut, common and kernel bunt, seed rots, and seedling diseases.
<i>carboxin</i> + <i>ipconazole</i> Rancona V100	Wheat, Barley, Oats, Rye	0.9 – 1.5 fl oz	For control of seedborne and soilborne fungi.
<i>carboxin</i> + <i>thiram</i> Vitavax 200 RTU-Vitavax-Thiram	Wheat, Barley, Oats, Triticale Wheat, Oats, Barley	2 oz 2–4 oz	Controls loose smut and stinking smut. Controls seedling blights. See label for specific rate for grains.
<i>carboxin</i> + <i>PCNB</i> + <i>metalaxyl</i> Prevail	Wheat, Oats, Barley	2.5–5 oz (wheat) 1.6–3.3 oz (oats)	Controls loose smut, common and kernel bunt, seed rots, and seedling diseases from Pythium and Rhizoctonia.
<i>difenoconazole</i> Dividend	Wheat	0.5–1 oz	Controls loose smut and stinking smut.
<i>difenoconazole</i> + <i>mefenoxam</i> Dividend XL RTA Dividend XL Dividend Extreme	Wheat Wheat Wheat	5–10 oz 1–2 oz 0.5–1 oz	Controls loose smut, stinking smut, and Pythium damping-off. Grower and commercially applied.
<i>fludioxonil</i> Maxim 4FS	Barley, Millet, Oats, Rye, Sorghum, Triticale, Wheat	0.08–0.16 fl oz	Controls Fusarium, Rhizoctonia, Helminthosporium, and weakly pathogenic fungi such as Aspergillus and Penicillium.
<i>ipconazole</i> Rancona 3.8 FS Rancona Apex Vortex	Wheat, Barley, Oats, Rye	0.051–0.085 fl oz (3.8 FS) 5–8.3 fl oz (Apex)	Controls loose smut, common and kernel bunt, seed rots, and seedling diseases.
<i>Ipconazole</i> + <i>metalaxyl</i> Rancona Pinnacle	Wheat, Barley, Oats, Rye	5.0–8.33 fl oz	Controls seed rot, damping off seed and soil borne fungi, loose smut, and seeding common and kernel bunt.
<i>mefenoxan</i> Apron XL Apron XL-LS	Wheat, Barley, Millet, Oats, Rye, Sorghum, Triticale	0.042–0.08	Controls Pythium damping-off. Does not control smuts.
<i>metalaxyl</i> Allegiance Sebring Dyna-shield Belmont	Wheat, Barley, Millet, Oats, Rye, Sorghum, Triticale	See label	Controls Pythium damping-off. Does not control smuts.

For information on CruiserMaxx Cereals (*thiamethoxam* + *mefenoxam* + *difenoconazole*), CruiserMaxx Vibrance Cereals (*sedaxane* + *thiamethoxam* + *mefenoxam* + *difenoconazole*), Cruiser Vibrance Quattro (*thiamethoxam* + *mefenoxam* + *difenoconazole* + *sedaxane* + *fludioxonil*), and Gaucho XT (*imidacloprid* + *metalaxyl* + *tebuconazole*), See the Insect Management Section of this guide. Commercial treatment of small grain seed is preferred, but a drill box treatment can be used with many formulations. Drill-box treatment may not give control equal to commercial treatment.

FUNGICIDE	CROP	RATE/100 LB SEED	REMARKS
<i>metalaxyl + metconazole + clothianidin</i> NipsIt SUITE	Wheat, Oats, Barley	5–7.5 fl oz	Controls common smut, flag smut, loose smut, seed decay fungi, Fusarium seed scab, Pythium seed rot, and seedling diseases. Early season Fusarium seedling dieback, early season Rhizoctonia root rot, and early season common rot.
<i>penflufen</i> Evergol Prime	Wheat, Oats, Barley	0.32 fl oz	Controls loose smut, common and kernel bunt, seed rots, and seedling diseases.
<i>prothioconazole + penflufen + metalaxyl</i> Evergol Energy	Wheat, Oats, Barley	1 fl oz	Controls loose smut, common and kernel bunt, seed rots, and seedling diseases along with early suppression of powdery mildew, rust, and glume/leaf blotch.
<i>sedaxane</i> Vibrance	Wheat, Barley, Oats, Rye, Triticale	0.08–0.16 fl oz	Controls loose smut, seed decay seedling blight and damping-off caused by Rhizoctonia solani.
<i>sedaxane + difenconazole + mefenoxam</i> Vibrance Extreme	Wheat, Barley, Oats, Rye, Triticale	2.8–5.6 fl oz	Controls smuts and bunts, general seed rot, seedling blight, root rot, and damping-off caused by seed or soilborne Fusarium spp or Rhizoctonia spp, Seedling blight, and root rot, and damping-off caused by Pythium spp, seed borne Septoria, Septoria leaf blotch, and Fusarium seed scab.
<i>sedaxane + difenconazole + fludioxonil + mefenoxam</i> Vibrance Quattro	Wheat, Barley, Oats, Rye, Triticale	5 fl oz	Controls smuts and bunts, general seed rot, seedling blight, root rot, and damping-off caused by seed or soilborne Fusarium spp or Rhizoctonia spp, Seedling blight and root rot, and damping-off caused by Pythium spp, seed borne Septoria, Septoria leaf blotch, and Fusarium seed scab.
<i>tebuconazole</i> Raxil (in various combinations with other fungicides)	Wheat, Oats, Barley	3.5–4.6 fl oz	Controls loose smut and stinking smut. Controls seedling blights. Commercially-applied and drill-box formulations available.
Thiram	Wheat, Barley, Rye	See label	Controls seedling blights. Does not control smuts. Can be used for drill-box treatment.
<i>triadimenol</i> Baytan 30 RTU Baytan-Thiram	Wheat, Barley, Oats, Rye All	0.75–1.5 oz 4.5–9 oz	Controls loose smut and stinking smut. Controls smuts and seedling blights.
<i>triticonazole + metconazole</i> Charter F	Wheat, Barley, Oats, Rye	5.4 fl oz	Controls loose smut, common and kernel bunt, seed rots, and seedling diseases.

For information on CruiserMaxx Cereals (*thiamethoxam + mefenoxam + difenconazole*), CruiserMaxx Vibrance Cereals (*sedaxane + thiamethoxam + mefenoxam + difenconazole*), Cruiser Vibrance Quattro (*thiamethoxam + mefenoxam + difenconazole + sedaxane + fludioxonil*), and Gaucho XT (*imidacloprid + metalaxyl + tebuconazole*), See the Insect Management Section of this guide. Commercial treatment of small grain seed is preferred, but a drill box treatment can be used with many formulations. Drill-box treatment may not give control equal to commercial treatment.

DISEASE	CHEMICAL	MOA	RATE PER ACRE	REI (Hours)	REMARKS AND PRECAUTIONS
Stagonospora Leaf and Glume Blotch, Leaf Rust, Stripe Rust, Powdery Mildew, Tan Spot	<i>azoxystrobin</i> Quadris, Equation, Satori	11	6.2–10.8 oz 4–12 fl oz	4 H	Apply after Feekes 6 but not later than Feekes 10.5. Do not harvest treated wheat for forage. A crop oil concentrate adjuvant may be added at 1% v/v to optimize efficacy.
	<i>azoxystrobin + cyproconazole</i> Azure Xtra	11 + 3	3.5–6.8 fl oz	12 H	Apply product at 3.5 oz/A in the spring at Feekes 5. Apply 5–6.8 fl oz/A between Feekes 8–10.5.1.
	<i>azoxystrobin + propiconazole</i> Quilt, QuiltXcel, Avaris	11 + 3	7–14 oz	12 H	Applications may be made no closer than a 14-day interval. Quilt and QuiltXcel can be applied up to Feekes growth stage 10.5. QuiltXcel has a higher rate of <i>azoxystrobin</i> . Low rates of Quilt and QuiltXcel are used for spring suppression of early season diseases, 10.5 fl oz and above are used for flag leaf protection and maximizing yield potential.
	<i>azoxystrobin + tebuconazole</i> Custodia	11 + 3	6.4–8.6 fl oz	12 H	Should be applied prior to disease development up to late head emergence (Feekes 10.5). Do not apply after this stage.
	<i>azoxystrobin + flutriafol</i> Topguard EQ	3 + 11	4.0–7.0 fl oz	12 H	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Do not apply past Feekes 10.54. An adjuvant may be added at recommended rates.
	<i>benzovindiflupyr + azoxystrobin + propiconazole</i> Trivapro	7 + 11 + 3	9.4–13.7 fl oz	12 H	For disease control on the flag leaf, apply from Feekes 8 (Zadoks 37) through Feekes 10 (Zadoks 45). Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Trivapro fungicide is applied when the flag leaf is 50% to fully emerged. Trivapro fungicide can be applied through full head emergence (Feekes growth stage 10.5.4).
	<i>cyproconazole</i> Alto	3	3–5.5 fl oz		
	<i>fluoxapyroxad + pyraclostrobin</i> Priaxor	7 + 11	4–8 fl oz	12 H	Apply no later than the beginning of flowering (Feekes 10.5 Zadok's 59). Maximum number of applications per season is 2.
	<i>fluoxapyroxad + pyraclostrobin + propiconazole</i> Nexicor	7 + 11 + 3	7–13 fl oz	12 H	For optimal disease control, begin applications of Nexicor prior to disease development. To maximize yield potential, it is important to protect the flag leaf. Apply Nexicor immediately after flag leaf emergence, no later than the beginning of flowering (Feekes 10.5, Zadok's 59).
	<i>fluoxastrobin</i> Evito	11	2–4 fl oz	12 H	For optimum results, begin applications preventatively and continue on a 14–21 day interval. Do not make more than 2 sequential applications. Apply prior to disease development from Feekes 5 (Zadok's 31) up to late head emergence at Feekes 10.5 (Zadok's 59).
	<i>fluoxastrobin + tebuconazole</i> Evito T	11 + 3	4–6 fl oz	12 H	Apply a maximum of 2 applications per season Apply no later than Feekes 10.5. For optimum results, apply the first application at shooting-pseudostem erected (approximately Feekes 5, Zadok's 31) and a second application no later than heading completed (Feekes 10.5, Zadok's 54).
	<i>fluoxastrobin + flutriafol</i> Fortix Preemptor SC	11 + 3	2–3 fl oz 4–6 fl oz	12 H	For early season control. Apply Fortix when flag leaf is 50% to fully emerged. Apply preventative when conditions for disease are favorable for development.
	<i>metconazole</i> Caramba	3	10–14 oz	12 H	Maximum number of applications per season is 2. Minimum time from application to harvest is 30 days.
<i>picoxystrobin</i> Approach	11	3–4 fl oz 6–12 fl oz	12 H	For early season preventive disease control. Begin applications of Approach prior to disease development and continue on a 7–14 day interval, depending on the targeted disease. Use higher rate and shorter interval when disease pressure is high.	

DISEASE	CHEMICAL	MOA	RATE PER ACRE	REI (Hours)	REMARKS AND PRECAUTIONS
Stagonospora Leaf and Glume Blotch, Leaf Rust, Stripe Rust, Powdery Mildew, Tan Spot (continued)	<i>picoxystrobin + cyproconazole</i> Approach Prima SC	11 + 3	3.4–6.8 fl oz	12 H	For early season preventive disease control. Begin applications of Approach Prima prior to disease development and continue on a 7–14-day interval, depending on the targeted disease. Use higher rate and shorter interval when disease pressure is high.
	<i>pydiflumetofen + propiconazole</i> Miravis Ace	7 + 3	13.7 fl oz	12 H	
	<i>propiconazole</i> Tilt, Propimax	3	4 oz	12 H	Tilt can be applied until heading stage (Feekes 10.5). Do not apply Tilt after this growth stage to avoid possible illegal residues.
	<i>propiconazole + trifloxystrobin</i> Stratego	3 + 11	10 oz	12 H	Do not apply more than 2 applications of Stratego per season. Do not apply after Feekes 10.5.
	<i>prothioconazole</i> Proline	3	4.3–5 fl oz	12 H	For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Proline. Up to 2 applications of Proline may be made per year.
	<i>prothioconazole + tebuconazole</i> Prosaro	3 + 3	6.5–8.2 fl oz	12 H	Begin applications of Prosaro preventively when conditions are favorable for disease development. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Prosaro.
	<i>prothioconazole + trifloxystrobin</i> Stratego YLD Delaro 325 SC	3 + 11	4 fl oz	12 H	Begin applications preventatively when conditions are favorable for disease development. Do not make more than 2 applications/season. Do not apply after Feekes 10.5. Do not apply within 35 days of harvest.
	<i>pyraclostrobin</i> Headline	11	6–9 oz	12 H	Apply no later than Feekes 10.5.
	<i>pyraclostrobin + metconazole</i> Twinline Multiva	11 + 3	7–9 fl oz	12 H	Do not apply more than 2 applications/season. Do not apply after Feekes 10.5.
	<i>tebuconazole</i> Folicur Several others products have <i>tebuconazole</i> as the active ingredient. Check label of specific products.	3	4 fl oz	12 H	Folicur is no longer manufactured (2009). No end-user restrictions for disease control. Use until supply is exhausted. Not labeled for powdery mildew control. For all <i>tebuconazole</i> products, a maximum of 4 fl oz/A/season may be applied.
<i>tebuconazole + trifloxystrobin</i> Absolute Maxx SC	3 + 11	3–5 fl oz	12 H	Begin applications preventatively when conditions are favorable for disease development. For optimum disease control apply 5 fl oz at flag leaf stage (Feekes 8–9). For early season suppression of Tan Spot, Leaf Blight and Powdery Mildew, apply at 3–4 oz. Do not apply more than 5 fl oz/season. Do not apply after Feekes growth stage 10.5.2. Do not apply within 35 days of harvest. Do not use with adjuvants.	

Economic yield response to control wheat diseases is most likely to occur in fields with yield potentials of more than 50 bu/A and varieties with fair to poor resistance. Always follow label instructions, recommendations and restrictions.

WHEAT DISEASE CONTROL

DISEASE	CHEMICAL	MOA	RATE PER ACRE	REI/PHI (Hours or Days)	EFFICACY
Fusarium Head Blight (SCAB)	metconazole 8.6% Caramba 0.75 SL	3	13.5–17	12 H/ 30 D	G
	propiconazole 41.8% Tilt 3.6 EC	3	4	12 H/ Apply at 50% flowering	P
	prothioconazole 41% Proline 480 SC	3	5–5.7	12 H/ 30 D	G
	pydiflumetofen + propiconazole Miravis Ace	7 + 3	13.7 fl oz	12 H/	G
	*tebuconazole 38.7% Folicur 3.6 F	3	4	12 H/ 30 D	F
	prothioconazole 19% + tebuconazole 19% Prosaro 421 SC	3 + 3	6.5–8.2	12 H/ 30 D	G

Efficacy categories: P—Poor; F—Fair; G—Good; VG—Very Good; E—Excellent. Timing of fungicide application is crucial for the control of FHB. Research indicates that products within the *triazole* class of fungicides are most effective if applied at early flowering (Feekes 10.5.1). *Strobilurin* fungicides are not recommended for management of FHB. *Strobilurin* fungicides can increase the DON content of FHB-infected grain.

* A maximum of 4 fl oz of *tebuconazole*-containing products may be applied per acre per crop season. Table modified from 2018 fungicide table produced by “The North Central Regional Committee on Management of Small Grain Diseases (NCERA-184).”

OAT (FOR GRAIN) DISEASE CONTROL

Alfredo Martinez-Espinoza, Extension Plant Pathologist

DISEASE	CHEMICAL	MOA	RATE PER ACRE	REI	REMARKS AND PRECAUTIONS
Rust, Leaf Spot, Powdery Mildew, Tan Spot, Septoria, Stagonospora Leaf and Glume Blotch	azoxystrobin Quadris Satori	11	9.2–12.0 fl oz	4 H	Apply after Feekes 6 but not later than Feekes 10.5. Do not harvest treated wheat for forage. A crop oil concentrate adjuvant may be added at 1.0% v/v to optimize efficacy.
	azoxystrobin + propiconazole Quilt, QuiltXcel, Avaris	11 + 3	7–14 fl oz 10.5 fl oz	12 H	Applications may be made no closer than a 14-day interval. Quilt and QuiltXcel can be applied up to Feekes growth stage 10.5. QuiltXcel has a higher rate of <i>azoxystrobin</i> . Low rates of Quilt and QuiltXcel are used for spring suppression of early season diseases. A rate of 10.5 fl oz and above are used for flag leaf protection and maximizing yield potential.
	benzovindiflupyr + azoxystrobin + propiconazole Trivapro SE	7 + 11 + 3	9.4–13.7 fl oz	12 H	For disease control on the flag leaf, apply from Feekes 8 (Zadoks 37) through Feekes 10 (Zadoks 45). Protecting the flag leaf is important for maximizing the potential yield.
	fluoxapyroxad + pyraclostrobin Priaxor Xemium	7 + 11	4–8 fl oz	12 H	Apply no later than the beginning of flowering (Feekes 10.5, Zadok's 59). Maximum number of applications per season = 2.

DISEASE	CHEMICAL	MOA	RATE PER ACRE	REI	REMARKS AND PRECAUTIONS
Rust, Leaf Spot, Powdery Mildew, Tan Spot, Septoria, Stagonospora Leaf and Glume Blotch (continued)	<i>fluoxapyroxad</i> + <i>pyraclostrobin</i> + <i>propiconazole</i> Nexicor Xemium	7 + 11 + 3	7–13 fl oz	12 H	For optimal disease control, begin applications of Nexicor prior to disease development. To maximize yield potential it is important to protect the flag leaf. Apply Nexicor immediately after flag leaf emergence, no later than the beginning of flowering (Feekes 10.5, Zadok's 59).
	<i>metconazole</i> Caramba	3	10–14 fl oz	12 H	Maximum number of applications per season = 2; Minimum time from application to harvest = 30 days.
	<i>picoxystrobin</i> Approach	11	2–4 fl oz 6–12 fl oz	12 H	For early season preventive disease control. Begin applications of Approach prior to disease development and continue on a 7- to 14-day interval, depending on the targeted disease. Use higher rate and shorter interval when disease pressure is high.
	<i>propiconazole</i> Tilt Propimax	3	2–4 fl oz 4 fl oz	12 H	Tilt can be applied until heading stage (Feekes 10.5). Do not apply Tilt after this growth stage to avoid possible illegal residues.
	<i>propiconazole</i> + <i>trifloxystrobin</i> Stratego	3 + 11	7 fl oz	12 H	Do not apply more than 2 applications of Stratego per season. Do not apply after Feekes 10.5.
	<i>prothioconazole</i> Proline	3	5–5.7 fl oz	12 H	For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Proline. Up to two applications of Proline can be made per year.
	<i>pydiflumetofen</i> + <i>propiconazole</i> Miravis Ace	7 + 3	13.7 fl oz	12 H	Apply between Feekes growth stage 10.3 (Zadoks 55) up to 10.5.4 (Zadoks 71).
	<i>pyraclostrobin</i> Headline	11	6–9 fl oz	12 H	Apply no later than Feekes 10.5.
	<i>pyraclostrobin</i> + <i>metconazole</i> Twinline	11 + 3	7–9 fl oz	12 H	Do not apply more than 2 applications per season. Do not apply after Feekes 10.5.
Fusarium Head Blight (FHB, Scab) ¹	<i>metconazole</i> Caramba	3	13.5–17.0 fl oz	12 H	PHI 30 days.
	<i>propiconazole</i> Tilt 3.6EC Propimax	3	4.0 fl oz	12 H	
	<i>prothioconazole</i> Proline 480 SC	3	5.0–5.7 fl oz	12 H	PHI 30 days.
	<i>pydiflumetofen</i> + <i>propiconazole</i> Miravis Ace	7 + 3	13.7 fl oz	12 H	Apply no later than Feekes 10.5.4 (Zadoks 71).

1. Timing of fungicide application is crucial for the control of FHB. Research indicates that products within the triazole class of fungicides are most effective if applied at early flowering (Feekes 10.5.1).

Strobilurin fungicides are not recommended for the management of FHB. Strobilurin fungicides can increase the DON content of FHB-infected grain.

This information is provided only as a guide. By law, it is the responsibility of the pesticide applicator to read and follow all current label directions. No endorsement is intended for any products listed, nor is criticism meant for products not listed. Always check the label before applying for the most current rates and restrictions.

SMALL GRAIN WEED CONTROL

A. Stanley Culpepper and Jenna Vance, Extension Weed Scientist

SMALL GRAIN

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
WHEAT: PREPLANT NO-TILL						
Emerged annual weeds, suppresses volunteer corn	<i>paraquat</i> 2 SL 3 SL	22	2–4 pt 1.7–2.7 pt	0.5–1 0.63–1	24 H/ N/A	U.S. EPA has restricted the use of <i>paraquat</i> to certified applicators ONLY and applicators must take a specialized training before use. Apply before crop emerges. Add nonionic surfactant at 2 pt/100 gal of spray or crop oil concentrate at 1 gal/100 gal of spray. Control of 12" corn at 1.5 pt/A is about 80% but may provide acceptable control until frost.
Emerged annual weeds, control or suppression of perennial weeds; use the full rate for ryegrass and radish	<i>glyphosate</i> 3.57 SL (3 lb ae) 4 SL (3 lb ae) 5 SL (3.7 lb ae) 5.5 SL (4.5 lb ae) 5.88 S (4.8 lb ae) 6 SL (5 lb ae)	9	32–48 fl oz 24–36 fl oz 23–34 fl oz 22–32 fl oz 20–30 fl oz 19–29 fl oz	0.75–1.13 ae	4 H/ N/A	Apply before crop emerges; suggest 3 or more days before emergence. Adjuvant recommendation varies by brand used. Cool temperatures, especially at night, may slow or even reduce control. For perennial weeds, rate can be increased but is specific by species, refer to the label. Suggest higher rates only be applied prior to planting. Application of this product may be repeated as needed up to a maximum of 6 lb ae per acre per year (5 qts of PowerMax 3). <i>Best program for ryegrass: spray glyphosate first and follow with paraquat about 5–7 days later.</i>
Controls most weeds; use full <i>glyphosate</i> rate for ryegrass	<i>glyphosate</i> + <i>2,4-D amine</i> (3.8 L)	9 + 4	see <i>glyphosate</i> + 12–16 fl oz	0.75–1.13 ae + 0.36–0.48	48 H/ N/A	Check brand of <i>2,4-D</i> used as some labels prohibit planting within 29 days of application. Research suggests plantback intervals of at least 24 days and 1 inch of rain between application and planting may be needed. Without required rainfall, serious injury can occur. <i>Quelex offers much less injury potential than 2,4-D.</i>
Summer and winter annual weeds including wild radish, henbit, chickweed; use full <i>glyphosate</i> rate for ryegrass	<i>glyphosate</i> + <i>thifensulfuron-methyl</i> + <i>tribenuron-methyl</i> (Harmony Extra SG with Total Sol) 50 SG	9 + 2 + 2	see <i>glyphosate</i> + 0.45–0.9 oz	0.75–1.13 ae + 0.0094–0.0187 + 0.0047–0.0094	12 H/ N/A	May be used as a burndown treatment prior to or shortly after planting, but prior to wheat emergence (suggest at least 3 days before emergence because of the <i>glyphosate</i>).
Volunteer Roundup Ready Corn and ryegrass; use full rates of both products for ryegrass	<i>glyphosate</i> + <i>clethodim</i> (Select) 2 EC (Select Max) 0.97 EC	9 + 1	see <i>glyphosate</i> + 4–8 fl oz 9–16 fl oz	0.75–1.13 ae + 0.06–0.13 0.06–0.13	24 H/ N/A	Do not plant for 30 days after applying clethodim. Rainfall after application and before planting of 0.5" or more advised. Corn < 12 inch: Select 4–6 oz; Select Max 9–12 oz. Corn 12–24 inch: Select 6–8 oz; Select Max 12–14 oz. Ryegrass 2–6 inch: Select 8 oz; Select Max 16 oz.
Residual wild radish, henbit, chickweed control and ryegrass suppression from Valor	<i>glyphosate</i> + <i>flumioxazin</i> (Valor EZ) (Valor SX)	9 + 14	see <i>glyphosate</i> + 1–2 fl oz 1–2 oz	0.75–1.13 ae + 0.032–0.064	12 H/ N/A	For Valor, a minimum of 30 days must pass and 1 inch of rainfall/irrigation must occur between application and planting. Significant injury is likely if required rainfall does not occur. On sands, a plant back interval of 40 days is suggested.
Common chickweed, C. geranium, henbit, horseweed, soybean, small wild radish; use full <i>glyphosate</i> rate for ryegrass	<i>glyphosate</i> + <i>halauxifen-methyl</i> + <i>florasulam</i> (Quelex) 0.2 WG	9 + 4 + 2	see <i>glyphosate</i> + 0.75 oz	0.75–1.13 + 0.0048 + 0.0047	12 H/ N/A	Apply as a preplant burndown treatment prior to, or shortly after planting prior to emergence; suggest at least 3 days before emergence because of the <i>glyphosate</i> . Weeds should be less than 4". Label requires addition of non-ionic surfactant or crop oil concentrate. An application of Quelex can be made for burndown and again in-crop, rotation to cotton and soybean is 3 months. UGA research shows excellent crop tolerance with Quelex.

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
WHEAT: PREEMERGENCE						
Annual ryegrass suppression and annual broadleaf weeds	<i>chlorsulfuron</i> + <i>metsulfuron-methyl</i> Finesse 75 WDG	2 + 2	0.5 oz	0.0195 + 0.0039	4 H/ 45 D	Plant seed at least 1" deep; may stunt wheat on sandy soils. Do not use where a later application of Osprey or PowerFlex is anticipated. Plant only STS soybeans 6 or more months after application. Crop injury may result if an organophosphate is used. See label for rotational restrictions. A lower rate may be used for weeds other than ryegrass, see label.
WHEAT: DELAYED PREEMERGENCE						
Ideal use is for residual control of ryegrass resistant to POST herbicides; must be activated prior to weed emergence	<i>pyroxasulfone</i> Zidua 85 WG Zidua 4.17 SC	15	0.7–1 oz 1.25–1.75 fl oz	0.037–0.053	12 H/ N/A	Suggest planting seed 0.75–1.25" deep. Label does not allow an application to broadcast seedlings. Seed must be uniformly covered without furrows to avoid injury. Apply Zidua when 80% of germinated wheat seeds have a shoot at least 0.5" long up through wheat spiking. Rate can be increased to 0.075 lb ai on medium texture soils, see label. Label restricts irrigation until wheat is emerged. Avoid application if a long period of rain is expected during wheat emergence. To minimize ryegrass resistance: <i>If treating a field with either Anthem Flex, Fierce, or Zidua this year, do not apply these products in that field next year in wheat.</i>
	<i>pyroxasulfone</i> + <i>carfentrazone</i> Anthem Flex 4SE	15 + 14	2.0–3.64 fl oz	0.06–0.11 + .004–0.008	12 H/ 7 D	Suggest planting seed 0.75–1.25" deep. Label does not allow an application to broadcast seedlings. Seeds must be uniformly covered without furrows to avoid injury. Apply 2.0 to 2.73 fl oz/A on coarse soils and 2.75 to 3.64 fl oz/A on medium-texture soils when 80% of germinated seeds have a shoot at least 0.5" long up through wheat spiking. To minimize ryegrass resistance: <i>If treating a field with either Anthem Flex, Fierce, or Zidua this year, do not apply these products in that field next year in wheat.</i>
WHEAT: POSTEMERGENCE SPIKE THROUGH EARLY POST (FOR USE IN FIELDS WITH RYEGRASS RESISTANT TO POST HERBICIDES)						
Ideal use is for residual control of ryegrass resistant to POST herbicides, helps on radish and henbit	<i>flufenacet</i> + <i>metribuzin</i> Axiom 68 DF	15 + 5	4–8 oz	0.13–0.27 + 0.03–0.067	12 H/ N/A	Plant seed at least 1" deep. Apply to wheat between spike and 2 leaf. Must be activated prior to weed emergence for effective weed control. Preemergence applications can cause severe injury on light soils. For most GA soils, < 6 oz/A of product is ideal. Heavy rains following application may cause wheat stunting. Rotations: soybean and corn 0 months; cotton 8 months; many other crops 12 months; many root crops 18 months.
Ideal use is for residual control of ryegrass resistant to POST herbicides; must be activated prior to weed emergence	<i>pyroxasulfone</i> Zidua 85 WG Zidua 4.17 SC	15	1–2.5 oz 1.75–4 fl oz	0.053–0.133	12 H/ N/A	Apply to wheat (drilled or broadcast) between spiking and 4 tiller. Lower rate on coarse soils and young wheat. <i>Sequential applications may be made as to not exceed 2.5 oz of the dry formulation or 4 fl oz of the liquid formulation.</i> To minimize ryegrass resistance: <i>If treating a field with either Anthem Flex, Fierce, or Zidua this year, do not apply these products in that field next year in wheat.</i>
	<i>pyroxasulfone</i> + <i>carfentrazone</i> Anthem Flex 4SE	15 + 14	2.0–3.64 fl oz	0.06–0.11 + 0.004–0.008	12 H/ 7 D	Apply 2.0 to 2.73 fl oz/A on coarse soils and 2.75 to 3.64 fl oz/A on medium-texture soils to wheat (drilled or broadcast) between spiking and 4 tiller. The label suggest the use of an adjuvant is recommended for consistent control; however, the use of this product is really for residual ryegrass activity. Sequential applications may be made not to exceed 4.55 oz/A . To minimize ryegrass resistance: <i>If treating a field with either Anthem Flex, Fierce, or Zidua this year, do not apply these products in that field next year in wheat.</i>

SMALL GRAIN WEED CONTROL

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
WHEAT: POSTEMERGENCE SPIKE THROUGH EARLY POST (FOR USE IN FIELDS WITH RYEGRASS RESISTANT TO POST HERBICIDES) (continued)						
Residual control of annual ryegrass, wild radish, and other weeds; must be activated prior to weeds reaching 0.25" for control	<i>pyrooxasulfone</i> + <i>flumioxazin</i> Fierce 76 WDG Fierce EZ 3.04 SC	15 +	1.5 oz 3 fl oz	0.04 +	12 H/ N/A	Drill seeds 1 to 1.25" deep; applications to heavy sands and low organic matter areas coupled with heavy rain after application may cause excessive injury. Application to fields where wheat seed has been broadcast and shallow incorporated will likely result in substantial crop damage. Apply to wheat from spike through the 2-leaf stage; DO NOT APPLY Preemergence. Apply only in water; no additives. Visual leaf tip burn and chlorosis is expected. Ideally, Fierce is activated after wheat is up but before weed emergence. No rotational concerns for corn, cotton, peanut, or soybean. To minimize ryegrass resistance: <i>If treating a field with either Anthem Flex, Fierce or Zidua this year, do not apply these products in that field next year in wheat.</i>
WHEAT: POSTEMERGENCE						
Likely not the best option for most GA fields; can be effective on cocklebur, nightshades, field pennycress, shepherdspurse, wild buckwheat, and morningglory	<i>bromoxynil</i> (Brox) 2 EC	5	1.5–2 pt	0.375–0.5	24 H/ 45 D	Apply to wheat from emergence to the boot stage. Apply to susceptible broadleaf weeds up to the 4 leaf stage, 2 inch height, or 1 inch diameter, whichever comes first. May be able to control 1–2" wild radish.
Common chickweed, C. geranium, henbit, horseweed, soybean, small wild radish	<i>halauxifen-methyl</i> + <i>florasulam</i> (Quelex) 0.2 WG	4 +	0.75 oz	0.0048 +	12 H/ 60 D	Apply to wheat between 2-leaf and flag leaf. Weeds should be less than 4"; stressed weeds may not be controlled. Mixtures with MCPA have shown improved wild radish control; be sure to follow MCPA application timing restrictions if applying the mixture. Add crop oil concentrate (0.5–1% v/v). Rotation of 3 months for cotton, corn, soybean and 9 months for peanut. See label about mixing with liquid nitrogen. UGA research shows excellent control of small radish but less effective on larger plants.
Non-resistant emerged annual ryegrass; small wild radish, henbit and chickweed Usually effective on annual bluegrass	<i>mesosulfuron-methyl</i> Osprey 4.5 WDG	2	4.75 oz	0.013	4 H/ 60 D	Apply to wheat between emergence and jointing to control ryegrass with less than 2 tillers. Add a nonionic surfactant (at least 80% active) at 2 qts/100 gal spray solution plus <i>ammonium nitrogen</i> fertilizer (28-0-0, 30-0-0, 32-0-0) at 1–2 qt/A. DO NOT topdress within 14 days of application or mix with 2,4-D or MCPA. Do not use liquid <i>nitrogen</i> as the carrier. May mix Osprey with Harmony Extra. Cotton/soybean can be planted 90 days after application. To minimize ryegrass resistance: <i>If treating a field with either Osprey or PowerFlex this year, do not apply either product on that field next year.</i>
Non-resistant emerged annual ryegrass	<i>pinoxaden</i> + <i>fenoxaprop-p-ethyl</i> Axial Bold 0.685 EC	1	15 fl oz	0.054 +	48 H/ 70 D	Apply to wheat between emergence and pre-boot to control ryegrass with less than 2 tillers. No adjuvant mentioned on label. Mixtures with Harmony Extra will offer broadleaf control. UGA data suggests not mixing with <i>nitrogen</i> but label allows water/ <i>nitrogen</i> mixtures containing up to 50% liquid <i>nitrogen</i> by volume. Add water to tank, then add Axial; then mix thoroughly and add <i>nitrogen</i> . One application per crop and any crop can be planted 90 days later. To minimize ryegrass resistance: <i>If treating a field with Axial this year, do not apply it on that field next year.</i>

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
WHEAT: POSTEMERGENCE (continued)						
Non-resistant emerged annual ryegrass and wild radish	<i>pyroxsulam</i> PowerFlex HL 13.13 WDG	2	2 oz	0.0164	12 H/ 60 D	Apply to wheat between 3 leaf and jointing to control ryegrass with less than 2 tillers. Add crop oil concentrate at 1–1.25% v/v (1–1.25 gal/ 100 gal spray solution). Can tank mix with Harmony Extra. UGA data suggests not mixing with <i>nitrogen</i> but label allows water- <i>nitrogen</i> mixture containing up to 50% liquid <i>nitrogen</i> by volume (< 30 lb/A of nitrogen). If applying in liquid <i>nitrogen</i> , use a nonionic surfactant at 0.25% v/v, instead of crop oil. An independent liquid <i>ammonium nitrogen</i> fertilizer application should not be made within 7 days of application; also, do not apply organophosphates within 5 days of PowerFlex. As a rotation crop, soybeans and cotton may be planted 3 months following application. To minimize ryegrass resistance: <i>If treating a field with either Osprey or PowerFlex this year, do not apply either product on that field next year.</i>
Fair residual control of annual ryegrass; if activated	<i>pendimethalin</i> Prowl H ₂ O 3.8 AS	3	1.5–2.5 pt	0.71–1.19	24 H/ 60 D	Apply to wheat after 1 leaf and before flag leaf. Prowl will not control emerged weeds and must be activated prior to weed emergence. May tank mix with any postemergence herbicide labeled for use in wheat. <i>Zidua is more effective on ryegrass but having both Zidua and Prowl in the system would improve resistance management.</i>
Wild garlic, curly dock, many other winter annual broadleaf weeds; will not control primrose and wild radish must be < 1"	<i>thifensulfuron-methyl</i> + <i>tribenuron-methyl</i> Harmony Extra SG with TotalSol 50 SG Harmony Extra, Nimble, others 75 WDG	2 + 2	0.45–0.9 oz 0.3–0.6 oz	0.0094–0.0187 + 0.0047–0.0094	12 H/ 45 D	Apply to wheat after 2 leaf and before flag leaf. Most winter annuals can be controlled with 0.75 oz/A of Harmony Extra 50SG; however, 0.75–0.9 oz/A is recommended for controlling wild garlic or very small wild radish. Apply to non-stressed weeds with less than 4 leaves when temperatures are above 50° F. Garlic should be less than 12" and should have 2–4" of new growth. Make no more than 2 applications per year applying a max of 1.5 oz/A of Harmony Extra Total Sol or equivalent active ingredient with other products. A nonionic surfactant at the rate of 1 qt/100 gal of spray solution is suggested when applied in water. Liquid <i>nitrogen</i> may be used as the carrier; in this case, premix the herbicide in water and add the mixture to <i>nitrogen</i> with agitation; add 0.5–1 pt nonionic surfactant to 100 gal spray solution if less than 50% of the total spray solution is liquid nitrogen. For radish , consider a tank mix with 0.375–0.5 lb active ingredient of <i>MCPA</i> or <i>2,4-D</i> (12–16 oz/A of 3.8 lb ai/A material). Add 0.5–1 pt nonionic surfactant/100 gal spray solution, surfactant rate can be increased to 1 qt but expect more injury. If mixing <i>2,4-D</i> or <i>MCPA</i> with Harmony Extra and using <i>nitrogen</i> as carrier, eliminate surfactant. Follow wheat stage of growth restrictions for <i>2,4-D</i> or <i>MCPA</i> .
Partial control of wild garlic, henbit and wild radish Harmony Extra is usually much more effective	<i>tribenuron-methyl</i> Express SG with TotalSol 50 SG Express 75 WDG	2	0.25–0.5 oz 0.167–0.33 oz	0.008–0.0156	12 H/ 45 D	Apply to wheat after 2 leaf and before flag leaf. Add 1 qt of nonionic surfactant/100 gal of spray solution. Consider mixtures with 0.375–0.5 lb active ingredient of <i>2,4-D</i> or <i>MCPA</i> for improved control of wild radish (add 1 pt nonionic surfactant/100 gal spray solution). If mixing <i>2,4-D</i> or <i>MCPA</i> with Express and using <i>nitrogen</i> as the carrier, use at most 0.5 pt of nonionic surfactant/100 gal of spray solution. Follow wheat stage of growth restrictions for <i>MCPA</i> or <i>2,4-D</i> when using these mixtures.

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

SMALL GRAIN WEED CONTROL

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
WHEAT: POSTEMERGENCE (continued)						
Most winter annual broadleaf weeds except chickweed, henbit, geranium, clover, red sorrel and knawel	<i>2,4-D amine</i> various brands 3.8 L	4	1–1.25 pt	0.48–0.6	48 H/ 14 D	Apply to fully tillered wheat only. Spraying wheat too young or after jointing may reduce yields. Better results obtained when daytime temperatures are above 50° F. For corn cockle, wild onion, and wild garlic rate can be increased to 2 pt/A (of 3.8 lb product) but crop injury is expected. Injury is increased when using liquid <i>nitrogen</i> as the carrier. Ester formulations can be added directly into <i>nitrogen</i> . If using amine formulation, premix in water (1 part <i>2,4-D</i> to 4 parts water) and add mixture to <i>nitrogen</i> with strong agitation. Amine formulations give less burn than ester formulations in <i>nitrogen</i> . Ester formulations may be more effective on weeds in very cold conditions. Amine formulations are suggested to minimize off-target movement. Consider mixtures with Harmony Extra or with Quelex. One application of <i>2,4-D</i> in-season only.
	<i>2,4-D ester</i> various brands 3.8 L		1–1.25 pt	0.48–0.6		
	<i>2,4-D ester</i> various brands 5.5 L		0.67–0.84 pt	0.48–0.6		
Most winter annual broadleaf weeds except chickweed, henbit, geranium, clover, red sorrel and knawel	<i>MCPA</i> various brands 3.7 SL	4	11–20 fl oz	0.347–0.58	48 H/ —	Research suggests to apply 12–16 oz/A when wheat has at least tillered and 16–20 oz/A when fully tillered but before boot stage. However, some labels limit the rate to 16 oz/A and the application timing to full tiller, check product label used closely. Safer on the crop than <i>2,4-D</i> ; slightly less effective on larger weeds when applied alone. No spray additive needed. Consider mixtures with Harmony Extra or with Quelex. Amine formulations are suggested to minimize off-target movement. MCPA will provide more residual activity on radish than <i>2,4-D</i> , although control may only be a week or so.
WHEAT: PRE-HARVEST						
Annual broadleaf and grass weeds, suppression of perennial weeds	<i>glyphosate</i> 3.57 SL (3 lb ae) 4 SL (3 lb ae) 5 SL (3.7 lb a.e) 5.5 SL (4.5 lb ae) 5.8 SL (4.88 lb ae) 6 SL (5 lb ae)	9	2.6 pt 2 pt 1.6 pt 22 fl oz 20 fl oz 20 fl oz	0.75 ae	4 H/ 7 D	Apply after hard dough stage of grain (30% or less grain moisture) but at least 7 days before harvest. Do not apply to wheat grown for seed. A wiper application could be used for only rope/sponge applicators (33–75% of solution with water) or a panel applicator (33–100% solution with water); do not use a roller applicator. Do not add a surfactant, and there is a 35 day pre-harvest interval for wiper applications.
Annual broadleaf weeds	<i>2,4-D amine</i> various brands 3.8 SL	4	1 pt	0.48	48 H/ 14 D	Apply when grain is in the hard dough stage (30% or less grain moisture) or later. Use only amine formulations to reduce volatility as sensitive crops are likely nearby during this time of year. DO NOT allow drift to any sensitive crop!
BARLEY: PREPLANT NO-TILL						
Emerged annual weeds, suppresses volunteer corn	<i>paraquat</i> 2 SL 3 SL	22	2–4 pt 1.7–2.7 pt	0.5–1 0.63–1	24 H/ N/A	U.S. EPA has restricted the use of <i>paraquat</i> to certified applicators ONLY and applicators must take a specialized training before use. Apply before crop emerges. Add nonionic surfactant at 2 pt/100 gal of solution or crop oil concentrate at 1 gal/100 gal of solution. Control of 12" corn at 1.5 pt/A is about 80% but may provide acceptable control until frost; see label.
Emerged annual weeds, control or suppression of perennial weeds; use full rate for ryegrass and radish	<i>glyphosate</i> 3.57 SL (3 lb ae) 4 SL (3 lb ae) 5 SL (3.7 lb ae) 5.5 SL (4.5 lb ae) 5.88 S (4.8 lb ae) 6 SL (5 lb ae)	9	32–48 fl oz 24–36 fl oz 23–34 fl oz 22–32 fl oz 20–30 fl oz 19–29 fl oz	0.75–1.13 ae	4 H/ N/A	Apply before crop emerges; suggest 3 or more days before emergence. Adjuvant recommendation varies by brand used. Cool temperatures, especially at night, may slow or even reduce control. For perennial weeds, rate can be increased but is specific by species, refer to the label. Suggest higher rates only be applied prior to planting. Application of this product may be repeated as needed up to a maximum of 6 lb ae per acre <u>per year</u> (5 qts of PowerMax 3). <i>Best program for ryegrass: spray glyphosate first and follow with paraquat about 5–7 days later.</i>

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
BARLEY: PREPLANT NO-TILL (continued)						
Control of most winter weeds; use full glyphosate rate for ryegrass	<i>glyphosate</i>	9	see <i>glyphosate</i>	0.75–1.13 ae	48 H/ N/A	Check brand of 2,4-D used as some labels prohibit planting within 29 days of application. Research suggests plantback intervals of at least 24 days and 1 inch of rain between application and planting may be needed. Without required rainfall, serious injury can occur.
	+ 2,4-D amine (3.8 SL)	4	+ 12–16 fl oz	+ 0.36–0.48		
Summer and winter annual weeds including wild radish, henbit, chickweed, ryegrass	<i>glyphosate</i>	9	see <i>glyphosate</i>	0.75–1.13 ae	12 H/ N/A	May be used as a burndown treatment prior to or shortly after planting, but prior to barley emergence (suggest at least 3 days before emergence because of the <i>glyphosate</i>). Use full <i>glyphosate</i> rate for ryegrass!
	+ <i>thifensulfuron-methyl</i>	2	+ 0.45–0.9 oz	+ 0.0094–0.0187		
	+ <i>tribenuron-methyl</i> (Harmony Extra SG with Total Sol) 50 SG	2		+ 0.0047–0.0094		
Volunteer Roundup Ready Corn and ryegrass; use full rates for ryegrass	<i>glyphosate</i>	9	see <i>glyphosate</i>	0.75–1.13	24 H/ N/A	Do not plant for 30 days after applying <i>clethodim</i> . Rainfall after application and before planting of 0.5" or more advised. Corn < 12 inch: Select 4–6 oz; Select Max 9–12 oz. Corn 12–24 inch: Select 6–8 oz; Select Max 12–16 oz. Ryegrass 2–6 inch: Select 8 oz; Select Max 14 oz
	+ <i>clethodim</i>	1	+ 4–8 fl oz 9–16 fl oz	+ 0.06–0.13 0.06–0.13		
	(Select) 2 EC (Select Max) 0.97 EC					
Common chickweed, C. geranium, henbit, horseweed, soybean, wild radish; use full <i>glyphosate</i> rate for ryegrass	<i>glyphosate</i>	9	see <i>glyphosate</i>	0.75–1.13	12 H/ N/A	Apply as a preplant burndown treatment prior to, or shortly after planting prior to emergence; suggest at least 3 days before emergence because of the <i>glyphosate</i> . Label requires addition of non-ionic surfactant or crop oil concentrate. An application of Quelex can be made for burndown and again in-crop, rotation to cotton and soybean is 3 months. UGA has not studied Quelex on barley.
	+ <i>halauxifen-methyl</i>	4	+ 0.0048	+ 0.0048		
	+ <i>florasulam</i>	2	+ 0.75 oz	+ 0.0047		
	(Quelex) 0.2 WG					
BARLEY: POSTEMERGENCE						
Likely not the best option for most GA fields, can be effective on cocklebur, nightshades, field pennycress, wild buckwheat, and morningglory	<i>bromoxynil</i> (Brox) 2 EC	5	1.5–2 pt	0.375–0.5	24 H/ 45 D	Apply on barley from emergence to the boot stage. Apply to susceptible broadleaf weeds up to the 4-leaf stage, 2 inch height, or 1 inch diameter, whichever comes first. May be able to control 1–2" wild radish.
Non-resistant emerged annual ryegrass	<i>pinoxaden</i> + <i>fenoxaprop-p-ethyl</i> Axial Bold 0.685 EC	1	15 fl oz	0.054 + 0.027	48 H/ 70 D	Apply to barley from emergence up to just before jointing to control ryegrass with less than 2 tillers. No adjuvant needed. Mixtures with Harmony Extra will offer broadleaf control. UGA suggests not mixing with <i>nitrogen</i> but label allows water/ <i>nitrogen</i> mixtures containing up to 50% liquid <i>nitrogen</i> by volume. Add water to tank, then add Axial; then mix thoroughly and add <i>nitrogen</i> . One application per crop and any crop can be planted 90 days later. To minimize ryegrass resistance: <i>If treating a field with Axial this year, do not apply it on that field next year.</i>
Common chickweed, C. geranium, henbit, horseweed, soybean, small wild radish	<i>halauxifen-methyl</i>	4	0.75 oz	0.0048	12 H/ 60 D	Apply to barley after 2 leaf and before flag leaf. Weeds should be less than 4" and not stressed. Add non-ionic surfactant (0.2–0.5% v/v) or crop oil concentrate (0.5 to 1% v/v). Rotation of 3 months for cotton, corn, and soybean, and 9 months for peanut. See label about mixing with liquid <i>nitrogen</i> . UGA has not studied Quelex on barley.
	+ <i>florasulam</i> (Quelex) 0.2 WG	2		+ 0.0047		

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

SMALL GRAIN WEED CONTROL

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
BARLEY: POSTEMERGENCE (continued)						
Wild garlic, curly dock, many other winter annual broadleaf weeds; will not control primrose and wild radish must be < 1"	<i>thifensulfuron-methyl</i> + <i>tribenuron-methyl</i>	2 + 2		0.0094–0.0187 + 0.0047–0.0094	12 H/ 45 D	Apply to barley after 2 leaf and before flag leaf. Most winter annuals can be controlled with 0.75 oz/A of Harmony Extra 50 SG with TotalSol; however, 0.75–0.9 oz/A is recommended for wild garlic or small wild radish. Add 1 qt of nonionic surfactant/100 gal of spray solution. Apply to non-stressed weeds with less than 4 leaves when temperatures are above 50°F. Garlic should be less than 12" tall with 2–4" of new growth. Liquid <i>nitrogen</i> may be used as the carrier. In this case, premix the herbicide in water and add the mixture to <i>nitrogen</i> with agitation; add 0.5–1 pt nonionic surfactant to 100 gal spray solution if less than 50% of the total spray solution is liquid nitrogen. For radish , consider a tank mix with 0.375–0.5 lb ai of <i>2,4-D</i> or <i>MCPA</i> (12–16 fl oz/A of 3.8 lb material). Do not use surfactant if applying with <i>2,4-D</i> or <i>MCPA</i> in <i>nitrogen</i> . Follow barley stage of growth restrictions for <i>2,4-D</i> or <i>MCPA</i> with mixtures.
	Harmony Extra SG with TotalSol 50 SG		0.45–0.9 oz			
	Harmony Extra, Nimble 75 WDG		0.3–0.6 oz			
Partial control of wild garlic, henbit and wild radish Harmony Extra is usually more effective	<i>tribenuron-methyl</i>	2		0.0078–0.0155	12 H/ 45 D	Apply to barley after 2 leaf and before flag leaf. Add 1 qt of nonionic surfactant/100 gal of spray solution. Consider mixing with 0.375–0.5 lb active ingredient of <i>MCPA</i> or <i>2,4-D</i> (12–16 fl oz of 3.8 lb ai material) for improved control of wild radish and other broadleaf weeds. If applying with <i>2,4-D</i> or <i>MCPA</i> , follow growth restrictions for these herbicides.
	Express SG TotalSol 50 SG Express 75 WDG		0.25–0.5 oz 0.167–0.33 oz			
Most winter annual broadleaf weeds except chickweed, henbit, geranium, red sorrel, clover and knawel	<i>MCPA</i> various brands 3.7 SL	4	12–20 fl oz	0.35–0.58	48 H/ —	Research suggests to apply 12–16 oz/A after tillering and 16–20 oz/A when fully tillered but before boot stage. However, some labels limit the rate to 16 oz/A and the application timing to full tiller, check product label used closely. Safer on crop than <i>2,4-D</i> ; slightly less effective on large weeds when applied alone. <i>Amine</i> formulation suggested to help minimize drift. No spray additive needed. Consider mixtures with Harmony Extra. <i>MCPA</i> will provide more residual activity on radish than <i>2,4-D</i> , although control may only be a week or so.
Most winter annual broadleaf weeds except chickweed, henbit, geranium, red sorrel, clover and knawel	<i>2,4-D amine</i> various brands 3.8 SL	4	1–1.25 pt	0.48–0.6	48 H/ 14 D	Apply to fully tillered barley only. Spraying barley too young or after jointing may reduce yields. Increase rate by 50% to control corn cockle. For corn cockle, wild onion, and wild garlic rate can be increased to 2 pt/A (of 3.8 lb product) but crop injury is expected. Better results are obtained when day-time temperatures are above 50°F. Liquid <i>nitrogen</i> may be used as a carrier for <i>2,4-D</i> . Ester formulations can be added directly into <i>nitrogen</i> . If using <i>amine</i> formulation, premix in water (1 part <i>2,4-D</i> to 4 parts water) and add mixture to <i>nitrogen</i> with strong agitation. <i>Amine</i> formulations give less burn than ester formulations in <i>nitrogen</i> while esters may perform slightly better in very cold conditions. Ester formulations are far more volatile and should be avoided if possible. Consider mixtures with Harmony Extra. One application in-season only.
	<i>2,4-D ester</i> various brands 3.8 SL		1–1.25 pt	0.48–0.6		
	<i>2,4-D ester</i> various brands 5.7 SL		0.67–0.84 pt	0.48–0.6		
BARLEY: PREHARVEST						
Annual broadleaf weeds	<i>2,4-D amine</i> various brands 3.8 SL	4	1 pt	0.48	48 H/ 14 D	Apply when grain is in hard dough stage or later. Do not allow drift to any sensitive crop! Apply only labeled AMINE formulations to reduce volatility.
Annual weeds, suppression of perennials	<i>glyphosate</i> Roundup PMAX3 5.8 SL (4.88 lb ae/A)	9	20 fl oz	0.75 ae	4 H/ 7 D	FEED BARLEY ONLY. Apply after the hard-dough stage and when the grain contains 20% moisture or less. Do not apply on barley grown for seed! Apply at least 7 days prior to harvest or grazing. A wiper application could be used for only rope/sponge applicators (33–75% of solution with water) or a panel applicator (33–100% solution with water); do not use a roller applicator. Do not add a surfactant and there is a 35 day pre-harvest interval for wiper applications.

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
OATS: PREPLANT NO-TILL						
Emerged annual weeds, control or suppression of perennial weeds; use the full rate for ryegrass and radish.	<i>glyphosate</i> 3.57 SL (3 lb ae) 4 SL (3 lb ae) 5 SL (3.7 lb ae) 5.5 SL (4.5 lb ae) 5.8 SL (4.88 lb ae) 6 SL (5 lb ae)	9	32–48 fl oz 24–36 fl oz 23–34 fl oz 22–32 fl oz 20–30 fl oz 19–29 fl oz	0.75–1.13 ae	4 H/ N/A	Apply before crop emerges; suggest 3 or more days before emergence. Adjuvant recommendation varies by brand used. Cool temperatures, especially at night, may slow or even reduce control. For perennial weeds, rate can be increased but is specific by species; refer to the label. Suggest higher rates only be applied prior to planting. Application of this product may be repeated as needed up to a maximum of 6 lb ae per acre <u>per year</u> (5 qts of PowerMax 3).
Control of most winter weeds; use full glyphosate rate for ryegrass	<i>glyphosate</i> + <i>2,4-D amine</i> (3.8 SL)	9 + 4	see <i>glyphosate</i> + 12–16 fl oz	0.75–1.13 ae + 0.36–0.48	48 H/ N/A	Check brand of <i>2,4-D</i> used as some labels prohibit planting within 29 days of application. Research suggests plantback intervals of 24 days and 1 inch of rain between application and planting may be needed. Without required rainfall, serious injury can occur.
Volunteer Roundup Ready Corn and ryegrass; use full rates for ryegrass	<i>glyphosate</i> + <i>clethodim</i> (Select) 2 EC (Select Max) 0.97 EC	9 + 1	see <i>glyphosate</i> + 4–8 fl oz 9–16 fl oz	0.75–1.13 ae + 0.063–0.13 0.06–0.13	24 H/ N/A	Do not plant for 30 days after applying <i>clethodim</i> . Rainfall after application and before planting of 0.5" or more advised. Corn < 12 inch: Select 4–6 oz; Select Max 9–12 oz. Corn 12–24 inch: Select 6–8 oz; Select Max 12–14 oz. Ryegrass 2–6 inch: Select 8 oz; Select Max 16 oz
OATS: POSTEMERGENCE						
Wild garlic, curly dock, many other winter annual broadleaf weeds; will not control primrose and wild radish must be < 1"	<i>thifensulfuron-methyl</i> + <i>tribenuron-methyl</i> Harmony Extra SG with TotalSol 50 SG Harmony Extra, Nimble 75 WDG	2 + 2	0.45–0.6 oz 0.3–0.4 oz	0.0094–0.0125 + 0.0047–0.0063	12 H/ 45 D	Apply to oats after 2 leaf and before flag leaf. Apply to non-stressed weeds with less than 4 leaves when temperatures are above 50F. Garlic should be less than 12" tall with 2–4" of new growth. Add 1 qt of nonionic surfactant/100 gal of spray solution. See label for varietal restrictions. Liquid nitrogen may be used as the carrier. In this case, premix the herbicide in water and add the mixture to nitrogen with agitation; add 0.5–1 pt nonionic surfactant to 100 gal spray solution if less than 50% of the total spray solution is liquid nitrogen (burn may still be noted). Consider mixing with 0.375–0.5 lb ai of 2,4-D or MCPA (12–16 fl oz/A of 3.8 L product) to control wild radish. Do not use surfactant if applying with <i>2,4-D</i> or <i>MCPA</i> in nitrogen. Mixtures with <i>MCPA</i> or <i>2,4-D</i> must be applied to fully tillered oats only.
Partial control of wild garlic, henbit and wild radish; Harmony Extra is usually more effective	<i>tribenuron-methyl</i> Express SG TotalSol 50 SG Express 75 WDG	2	0.2 oz 0.133 oz	0.0078–0.0155	12 H/ 45 D	Apply to oats after 2 leaf but before flag leaf is visible. Add 1 qt of nonionic surfactant/100 gal of spray solution. Must be tank mixed with another registered herbicide; consider mixing with 0.375–0.5 lb active ingredient of <i>MCPA</i> or <i>2,4-D</i> (12–16 fl oz of 3.8 lb ai material) for improved control of wild radish and other broadleaf weeds. If applying with <i>2,4-D</i> or <i>MCPA</i> , follow growth restrictions for these herbicides.
Most winter annual broadleaf weeds except chickweed, henbit, geranium, red sorrel, clover and knawel	<i>MCPA</i> various brands 3.7 SL	4	12–16 fl oz	0.347–0.46	48 H/ —	Apply up to 16 fl oz/A when oat is full tiller; use higher rates only when fully tillered. Safer on oats than <i>2,4-D</i> ; slightly less effective on larger weeds when applied alone. No spray additive needed. Consider mixtures of MCPA with Harmony Extra for improved weed control.
Most winter annual broadleaf weeds except chickweed, henbit, geranium, red sorrel, clover and knawel	<i>2,4-D amine</i> various brands 3.8 SL	4	12–20 fl oz	0.36–0.59	48 H/ 14 D	Apply to fully tillered oats only. Spraying oats too young or after jointing may reduce yields. Oats are less tolerant of <i>2,4-D</i> than wheat. Better results are obtained when day-time temperatures are above 50°F. Liquid <i>nitrogen</i> may be used as a carrier for <i>2,4-D</i> . Premix in water (1 part <i>2,4-D</i> to 4 parts water) and add mixture to <i>nitrogen</i> with strong agitation. Notice only an <i>amine</i> formulation of <i>2,4-D</i> is recommended due to crop response. Consider mixtures of 2,4-D with Harmony Extra for improved weed control. One application in-season only.

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

SMALL GRAIN WEED CONTROL

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
OATS: PREHARVEST						
Annual broadleaf weeds	<i>2,4-D amine</i> various brands 3.8 SL	4	1 pt	0.48	48 H/ 14 D	Apply when grain is in hard dough stage or later. Do not allow drift to any sensitive crops. Apply only labeled AMINE formulations to reduce volatility and off-target movement.
RYE: PREPLANT						
Emerged annual weeds, control or suppression of perennial weeds; use the full rate for ryegrass and radish.	<i>glyphosate</i> 3.57 SL (3 lb ae) 4 SL (3 lb ae) 5 SL (3.7 lb a.e) 5.5 SL (4.5 lb ae) 5.8 SL (4.88 lb ae) 6 SL (5 lb ae)	9	32–48 fl oz 24–36 fl oz 23–34 fl oz 22–32 fl oz 20–30 fl oz 19–29 fl oz	0.75–1.13 ae	4 H/ N/A	Apply before crop emerges; suggest 3 or more days before emergence. Adjuvant recommendation varies by brand used. Cool temperatures, especially at night, may slow or even reduce control. For perennial weeds, rate can be increased but is specific by species, refer to the label. Suggest higher rates only be applied prior to planting. Application of this product may be repeated as needed up to a maximum of 6 lb ae per acre <u>per year</u> (5 qts of PowerMax 3).
Control of most winter weeds; use full <i>glyphosate</i> rate for ryegrass	<i>glyphosate</i> + <i>2,4-D amine</i> (3.8 SL)	9 + 4	see <i>glyphosate</i> + 12–16 fl oz	0.75–1.13 ae + 0.36–0.48	48 H/ N/A	Check brand of <i>2,4-D</i> used as some labels prohibit planting within 29 days of application. Research suggests plantback intervals of at least 24 days and 1 inch of rain between application and planting may be needed. Without required rainfall, serious injury can occur.
Volunteer Roundup Ready Corn and ryegrass; use full rates for ryegrass	<i>glyphosate</i> + <i>clethodim</i> (Select) 2 EC (Select Max) 0.97 EC	9 + 1	see <i>glyphosate</i> + 4–8 fl oz 8–16 fl oz	0.75–1.13 ae + 0.063–0.13 0.06–0.13	24 H/ N/A	Do not plant for 30 days after applying <i>clethodim</i> . Rainfall after application and before planting of 0.5" or more advised. Corn < 12 inch: Select 4–6 oz; Select Max 8–12 oz. Corn 12–24 inch: Select 6–8 oz; Select Max 12–16 oz. Ryegrass 2–6 inch: Select 8 oz; Select Max 16 oz.
RYE: POSTEMERGENCE						
Most winter annual broadleaf weeds except chickweed, henbit, geranium, red sorrel, clover and knawel	<i>MCPA</i> various brands 3.7 SL	4	12–20 fl oz	0.347–0.58	48 H/	Research suggests to apply 12–16 oz/A after tillering and 16–20 oz/A when fully tillered but before boot stage. However, some labels limit the rate to 16 oz/A and the application timing to full tiller; check product label used closely. Safer than <i>2,4-D</i> on crop; slightly less effective on larger weeds when applied alone. No spray additive needed. <i>MCPA</i> will provide more residual activity on radish than <i>2,4-D</i> , although control may only be a week or so.
	<i>2,4-D amine</i> various brands 3.8 SL	4	1–1.25 pt	0.48–0.6	48 H/ 14 D	Apply to fully tillered rye only. Spraying rye too young or after jointing can reduce yields. For corn cockle, wild onion, and wild garlic rate can be increased to 2 pt/A (of 3.8 lb product) but crop injury is expected. Better results are obtained when day-time temperatures are above 50°F. Liquid <i>nitrogen</i> may be used as a carrier for <i>2,4-D</i> . Ester formulations can be added directly into <i>nitrogen</i> . If using amine formulation, premix in water (1 part <i>2,4-D</i> to 4 parts water) and add mixture to <i>nitrogen</i> with strong agitation. Amine formulations give less burn than ester formulations in <i>nitrogen</i> . Ester may be more effective in cold conditions. Only in-season application only.
	<i>2,4-D ester</i> various brands 3.8 SL <i>2,4-D ester</i> various brands 5.7 SL		1–1.25 pt 0.67–0.84 pt			
Likely not the best option for most GA fields; can be effective on cocklebur, nightshades, field pennycress, shepherdspurse, wild buckwheat, and morningglory	<i>bromoxynil</i> (Brox) 2 EC	5	1.5–2 pt	0.375–0.5	24 H/ 45 D	Apply to rye from emergence to the boot stage. Apply to susceptible broadleaf weeds up to the 4 leaf stage, 2 inch height, or 1 inch diameter, whichever comes first. May be able to control 1–2" wild radish.

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
RYE: PREHARVEST						
Annual broadleaf weeds	<i>2,4-D amine</i> various brands 3.8 SL	4	1 pt	0.48	48 H/ 14 D	Apply when grain is in the hard dough stage or later. Do not allow drift to sensitive crops, especially cotton and tobacco. Apply only labeled AMINE formulations during this time of year.
TRITICALE: PREPLANT						
Emerged annual weeds, control or suppression of perennials; use the full rate for ryegrass and radish.	<i>glyphosate</i> 3.57 SL (3 lb ae) 4 SL (3 lb ae) 5 SL (3.7 lb ae) 5.5 SL (4.5 lb ae) 5.8 SL (4.88 lb ae) 6 SL (5 lb ae)	9	32–48 fl oz 24–36 fl oz 23–34 fl oz 22–32 fl oz 20–30 fl oz 19–29 fl oz	0.75–1.13 ae	4 H/ N/A	Apply before crop emerges; suggest 3 or more days before emergence. Adjuvant recommendation varies by brand used. Cool temperatures, especially at night, may slow or even reduce control. For perennial weeds, rate can be increased but is specific by species, refer to the label. Suggest higher rates only be applied prior to planting. Application of this product may be repeated as needed up to a maximum of 6 lb ae per acre <u>per year</u> (5 qts of PowerMax 3).
Summer and winter annual weeds including wild radish, henbit, chickweed; use full <i>glyphosate</i> rate for ryegrass	<i>glyphosate</i> + <i>thifensulfuron-methyl</i> + <i>tribenuron-methyl</i> (Harmony Extra SG with Total Sol) 50 SG	9 + 2 + 2	see <i>glyphosate</i> + 0.45–0.9 oz	0.75–1.13 ae + 0.0094–0.0187 + 0.0047–0.0094	12 H/ N/A	May be used as a burndown treatment prior to or shortly after planting, but prior to emergence (suggest at least 3 days before emergence because of the <i>glyphosate</i>).
Volunteer Roundup Ready Corn and ryegrass; use full rates for ryegrass	<i>glyphosate</i> + <i>clethodim</i> (Select) 2 EC (Select Max) 0.97 EC	9 + 1	see <i>glyphosate</i> + 4–8 fl oz 8–16 fl oz	0.75–1.13 ae + 0.063–0.13 0.065–0.13	24 H/ N/A	Do not plant for 30 days after applying <i>clethodim</i> . Rainfall after application and before planting of 0.5" or more advised. Corn < 12 inch: Select 4–6 oz; Select Max 8–12 oz. Corn 12–24 inch: Select 6–8 oz; Select Max 12–16 oz. Ryegrass 2–6 inch: Select 8 oz; Select Max 16 oz.
Common chickweed, <i>C. geranium</i> , henbit, horseweed, soybean, wild radish; use full <i>glyphosate</i> rate for ryegrass	<i>glyphosate</i> + <i>halauxifen-methyl</i> + <i>florasulam</i> (Quelex) 0.2 WG	9 + 4 + 2	see <i>glyphosate</i> + 0.75 oz	0.75–1.13 ae + 0.0048 + 0.0047	12 H/ N/A	Apply as a preplant burndown treatment prior to, or shortly after planting prior to emergence; suggest at least 3 days before emergence. Label requires addition of non-ionic surfactant or crop oil concentrate. An application can be made for burndown and again in-crop, rotation to cotton and soybean is 3 months. UGA has not studied Quelex on triticale.
TRITICALE: POSTEMERGENCE SPIKE THROUGH EARLY POST						
Ideal use is for residual control of ryegrass resistant to POST herbicides; helps on radish and henbit	<i>flufenacet</i> + <i>metribuzin</i> Axiom 68 DF	15 + 5	4–8 oz	0.136–0.027 + 0.34–0.068	12 H/ N/A	Triticale seed should be planted at least 1" deep. Apply to triticale between spike and 2 leaf. Must be activated prior to weed emergence for effective weed control. Preemergence applications can cause severe injury on light soils. For most Georgia soils, < 6 oz/A of product is ideal. Heavy rains following application may cause stunting. Avoid application if wet conditions after application are expected for a week. Rotation to soybean and corn is 0 months, cotton 8 months, many other crops 12 months and root crops 18 months. UGA has not studied Axiom on triticale.

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

SMALL GRAIN WEED CONTROL

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
TRITICALE: POSTEMERGENCE						
Most winter annual broadleaf weeds except chickweed, henbit, geranium, red sorrel, clover and knawel	<i>2,4-D amine</i> various brands 3.8 SL	4	12–20 oz	0.36–0.59	48 H/ 14 D	Apply to fully tillered triticale only. Spraying when too young or after jointing can reduce seed yields. For corn cockle, wild onion, and wild garlic rate can be increased to 2 pt/A (of 3.8 lb product) but crop injury is expected. Better results are obtained when day-time temperatures are above 50°F. Liquid nitrogen may be used as a carrier for <i>2,4-D</i> . If using amine formulation, premix in water (1 part <i>2,4-D</i> to 4 parts water) and add mixture to nitrogen with strong agitation. Only 1 application of <i>2,4-D</i> in crop allowed. Consider mixtures with Harmony Extra for improved control.
Likely not the best option for most GA fields, can be effective on cocklebur, nightshades, field pennycress, shepherdspurse, wild buckwheat, and morningglory	<i>bromoxynil</i> (Brox) 2 EC	5	1.5–2 pt	0.375–0.5	24 H/ 45 D	Apply to triticale between emergence and boot stage. Apply to susceptible broadleaf weeds up to the 4 leaf stage, 2 inch height, or 1 inch diameter, whichever comes first. May be able to control 1–2 inch wild radish.
Annual broadleaf weeds and ryegrass suppression	<i>chlorsulfuron</i> + <i>metsulfuron methyl</i> Finesse 75 WDG	2 + 2	0.2–0.4 oz	0.008 + 0.0016 to 0.016 + 0.0031	4 H/ 45 D	Apply to triticale after 2 leaf and before flag leaf. See label for weeds controlled, application rates, and use of surfactant. Plant only STS soybeans 6 months or more after application. Early application necessary for ryegrass suppression. See comments under wheat.
Common chickweed, C. geranium, henbit, horseweed, soybean, small wild radish	<i>halauxifen-methyl</i> + <i>florasulam</i> (Quelex) 0.2 WG	4 + 2	0.75 oz	0.0048 + 0.0047	12H/ 60 D	Apply to triticale between 2 leaf and flag leaf. Weeds should be less than 4" and not stressed. Add nonionic surfactant (0.2–0.5% v/v) or crop oil concentrate (0.5–1% v/v). Rotation of 3 months for cotton, corn, and soybean, and 9 months for peanut. See label about mixing with liquid <i>nitrogen</i> . UGA has not studied Quelex in triticale.
Non-resistant emerged annual ryegrass, small wild radish, henbit, and chickweed Usually effective on annual bluegrass	<i>mesosulfuron-methyl</i> Osprey 4.5 WDG	2	4.75 oz	0.013	4 H/ 60 D	For fall sown crop only. Apply to triticale between emergence and jointing to control ryegrass with less than 2 tillers. Add a nonionic surfactant (at least 80% active) at 2 qts/100 gal spray solution plus <i>ammonium nitrogen</i> fertilizer (28-0-0, 30-0-0, 32-0-0) at 1–2 qt/A. DO NOT topdress within 14 days of application. Do not use liquid nitrogen as the carrier. May mix with Harmony Extra. Cotton/soybean can be planted 90 days after application. To minimize ryegrass resistance: <i>If treating a field with either Osprey or PowerFlex this year, do not apply either product on that field next year.</i>
Non-resistant emerged annual ryegrass, also very effective on wild radish and several other broadleaf weeds	<i>pyroxsulam</i> PowerFlex HL 13.13 WDG	2	2 oz	0.0164	12 H/ 60 D	Apply to triticale between 3 leaf and jointing to control ryegrass with less than 2 tillers. Add crop oil concentrate at 1–1.25% v/v (1–1.25 gal/100 gal spray solution). May tank mix with Harmony Extra. UGA suggests not mixing with <i>nitrogen</i> but label allows water- <i>nitrogen</i> mixture containing up to 50% liquid <i>nitrogen</i> by volume (< 30 lb/A of nitrogen). If applying in liquid <i>nitrogen</i> , use a nonionic surfactant at 0.25% v/v, instead of crop oil. An independent liquid <i>ammonium nitrogen</i> fertilizer application should not be made within 7 days of application; also do not apply organophosphates within 5 days of PowerFlex. As a rotation crop, soybeans and cotton may be planted 3 months following application. To minimize ryegrass resistance: <i>If treating a field with either Osprey or PowerFlex this year, do not apply either product on that field next year.</i>

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

WEEDS CONTROLLED	HERBICIDE	MOA	AMOUNT OF FORMULATION (Broadcast Rate/Acre)	LBS ACTIVE INGREDIENT (Broadcast Rate/Acre)	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS (READ ALL LABELS)
TRITICALE: POSTEMERGENCE (continued)						
Fair residual control of annual ryegrass; if activated	<i>pendimethalin</i> Prowl H ₂ O 3.8 AS	3	1.5–2.5 pt	0.71–1.19	24 H/ 60 D	Apply to triticale after 1 leaf and before flag leaf. Prowl will not control emerged weeds and must be activated prior to weed emergence. May tank mix with any postemergence herbicide labeled for use in triticale.
Wild garlic, curly dock, many other winter annual broadleaf weeds; will not control primrose and wild radish should be < 1"	<i>thifensulfuron-methyl</i> +	2 +	0.45–0.9 oz	0.0094–0.0187 +	12 H/ 45 D	Apply to triticale between 2 leaf and flag leaf. Most winter annuals can be controlled with 0.75 oz/A of Harmony Extra 50 SG; however, 0.75–0.9 oz/A is recommended for controlling wild garlic or small wild radish. Add 1 qt of nonionic surfactant/100 gal of spray solution. Apply to non-stressed weeds with less than 4 leaves when temperatures are above 50° F. Garlic should be less than 12" tall and should have 2–4" of new growth. Liquid <i>nitrogen</i> may be used as the carrier. When using <i>nitrogen</i> as the carrier, add 0.5–1 pt nonionic surfactant to 100 gal spray solution if less than 50% of the total spray solution is liquid nitrogen (burn may still be noted). Tank mix with 0.375–0.5 lb active ingredient of 2,4-D (12–16 fl oz/A of 3.8 L product) for control of wild radish. Do not use surfactant if applying with 2,4-D in <i>nitrogen</i> . Mixtures with 2,4-D must be applied to fully tillered triticale only.
	<i>tribenuron-methyl</i> Harmony Extra SG with TotalSol 50 SG Harmony Extra, Nimble 75 WDG	2		0.0047–0.0094		
Partial control of wild garlic, henbit and wild radish Harmony Extra is usually more effective	<i>tribenuron-methyl</i> Express SG TotalSol 50 SG Express 75 WDG	2	0.25–0.5 oz 0.167–0.33 oz	0.0078–0.0155	12 H/ 45 D	Apply to triticale between 2 leaf and flag leaf. Add 1 qt of nonionic surfactant/100 gal of spray solution. Consider mixing with 0.375–0.5 lb active ingredient of 2,4-D (12–16 fl oz of 3.8 lb ai material) for improved control of wild radish and other broadleaf weeds. If applying with 2,4-D, follow growth restrictions for these herbicides.
TRITICALE: PREHARVEST						
Annual broadleaf weeds	2,4-D amine various brands 3.8 SL	4	1 pt	0.48	48 H/ 14 D	Apply when grain is in the hard dough stage or later. Do not allow drift to ANY sensitive crop. Apply only labeled AMINE formulations to reduce volatility.

Mode of Action (MOA) code can be used to delay weed resistance by increasing herbicide diversity in a management program.

SMALL GRAIN WEED RESPONSE TO HERBICIDES

A. Stanley Culpepper, Extension Weed Scientist

SMALL GRAIN

WEEDS	2,4-D ¹	MCPA ¹	EXPRESS ¹	QUELEX ¹	EXPRESS + MCPA ¹ OR 2,4-D	BUCTRIL ¹	HARMONY EXTRA ¹	HARMONY EXTRA + MCPA OR 2,4-D ¹	PEAK ¹	FINESSE ²
annual bluegrass	N	N	N		N	N	N	N	N	N
annual ryegrass	N	N	N	N	N	N	N	N	N	F
buttercup	G			F-G			G	G-E		G
common chickweed	P	P	G	G-E	G-E	P-F	G	G-E		G
common ragweed	G	F				E	P-F	F-G	E	
cornflower	G					G-E	P	F-G		F
cudweed	G-E	G-E			E	G	E	E		
curly dock	P	P			P	P-F	E	E		
dandelion	E	E			E	E		E		
dogfennel	G	F				G-E	E	E		
evening primrose	E	E		G-E	E	F-G	P-F	E	F-G	
field pennycress	G			G		G	G	G-E		G
geranium	F	F		G			F-G	G-E		
goldenrod	F	G				F				
hairy vetch	F	F				F	F-G	G-E		
henbit	P	P	F	G-E	G	F	G	G-E	F-G	G
horsenettle	F	F				F				
horseweed	F-G	F				F	F-G	F-G		
knawel	P					P	G	G		
lambsquarters	G	G				E	E	E	G	
plantains	E	E			E	E	E	E		
shepherd's-purse	G-E	G-E		G	E	G	E	E	G	G
swinecress	G	G			G-E	G-E	E	E		
thistles	G	G				G	F-G	G	F-G	
vetch	G			G-E		F	P			
Virginia pepperweed	E				E	F-G	G	E		
wild garlic	F	P				P	G-E	G-E	E	P
wild mustard	E	G-E	F	G-E ³	E	F-G	F-G	E	G	G
wild radish	E	G-E	F	G-E ³	E	F-G	F-G ³	E	G	G

E—excellent control, 90% or better
 G—good control, 80–90%
 F—fair control, 60–80%
 P—poor control, 30–60%
 N—no control, less than 30%

1. Timely postemergence application.
2. Applied preemergence.
3. Must be less than 2 inches.

WEEDS	AXIAL XL ¹	AXIOM ²	ZIDUA ²	FIERCE ²	OSPREY ¹	POWERFLEX ¹
annual bluegrass	N	G			G-E	P-F
annual ryegrass	G-E ³	P-G ⁴	G-E ⁵	G-E ⁵	G-E ⁶	G-E ⁶
buttercup	N					
common chickweed	N			G-E	F-G ⁷	F-G ⁷
common ragweed	N					
cornflower	N				P	
cudweed	N					
curly dock	N				P	
dandelion	N					
dogfennel	N					
evening primrose	N		P	E	P	P
field pennycress	N					
geranium	N	G		G		
goldenrod	N					
hairy vetch	N					
henbit	N	G-E	P-F	G-E	G-E	G
horsenettle	N					
horseweed	N			G-E		
knawel	N					
lambsquarters	N			E		
plantains	N					
shepherd's-purse	N					
swinecress	N		P	G-E	E	
thistles	N					
vetch	N				P-F ⁷	
Virginia pepperweed	N					
wild garlic	N				P	
wild mustard	N	G-E	P-F	G-E	G	G-E
wild radish	N	G-E	P-F	G-E	G	G-E

E—excellent control, 90% or better
 G—good control, 80–90%
 F—fair control, 60–80%
 P—poor control, 30–60%
 N—no control, less than 30%

1. Timely postemergence application.
2. Applied spike to wheat but PRE to weeds.
3. Axial and Hoelon have a similar mode of action. Axial controls about 85% of the Hoelon-resistant populations studied in GA.
4. Provides good control if Axiom is activated prior to ryegrass germination. Poor control if ryegrass emerges prior to Axiom activation.
5. For this level of control, Zidua must be activated prior to ryegrass emergence. For Fierce, ryegrass must be less than 0.25" when activated.
6. Will not control ALS-resistant ryegrass.
7. Weeds must not be larger than 2" at time of application.